

A CONJECTURE ABOUT ARTIN–TITS GROUPS

PATRICK DEHORNOY AND EDDY GODELLE

ABSTRACT. We conjecture that the word problem of Artin–Tits groups can be solved without introducing trivial factors ss^{-1} or $s^{-1}s$. Here we make this statement precise and explain how it can be seen as a weak form of hyperbolicity. We prove the conjecture in the case of Artin–Tits groups of type FC, and we discuss various possible approaches for further extensions, in particular a syntactic argument that works at least in the right-angled case.

Artin–Tits groups, also known as Artin groups, are those groups that are defined by relations of the form

$$(1) \quad sts\dots = tst\dots$$

where both terms consist of two alternating letters and have the same length. These groups were first investigated by J. Tits in the late 1960’s [4]. Two seminal references are [5] and [15]. At the moment many questions involving these groups remain open, including the decidability of the word problem in the general case [8].

The aim of this paper is to discuss a conjecture about the word problem of Artin–Tits groups, namely that, when a word w represents 1, hence is equivalent to the empty word modulo the relations, then w can be transformed to the empty word without introducing trivial factors ss^{-1} or $s^{-1}s$ at any intermediate step of the derivation. Here we shall elaborate this intuitive idea into a precise statement, called Property H, and interpret the latter as a weak form of hyperbolicity (whence the notation). So, our conjecture is that all Artin–Tits groups—or, rather, all Artin–Tits presentations since Property H involves a specific presentation—satisfy Property H. It is known that Artin–Tits groups are not hyperbolic in general and, therefore, our conjecture is about how much of hyperbolicity could nevertheless be true in these groups. The main positive result we prove here is that all Artin–Tits presentation of type FC satisfy Property H. In terms of applications, Property H does not imply the decidability of the word problem, at least directly, but it implies that the monoid defined by a family of relations embeds in the corresponding group. Such a result is known to be true for Artin–Tits groups [22], but, so far, the only available proof is quite indirect and relies on the existence of linear representations. If our conjecture is true, it would in particular lead to a more direct proof of that result.

The article is organized as follows. In Section 1, we introduce Property H, connect it with hyperbolicity, and show as a warm-up exercise that all Artin–Tits presentations of spherical type satisfy Property H. We also mention the application to monoid embeddability. Next, in Section 2, we establish that all Artin–Tits presentations of type FC satisfy Property H. Finally, in Section 3, we give an alternative proof of the result that all right-angled Artin–Tits presentations (a subfamily of type FC) satisfy Property H. The current range of the

1991 *Mathematics Subject Classification.* 20B30, 20F55, 20F36.

Key words and phrases. Artin–Tits groups, positive presentation, van Kampen diagram, Dehn algorithm, amalgamated product, word problem, subword reversing.

Work partially supported by the ANR grant TheoGar ANR-08-BLAN-0269-02.

argument is more limited than that of Section 2, but it might be more promising in terms of subsequent developments as it involves no normal form.

1. PROPERTY H

We introduce special transformations and Property H in Subsection 1.1. We discuss its connection with the Dehn algorithm in Subsection 1.2. Then, in Subsection 1.3, we establish that all Artin–Tits presentations of spherical type satisfy Property H. Finally, we mention in Subsection 1.4 an application of Property H to the embeddability of a monoid $\langle S|R \rangle^+$ in the corresponding group $\langle S|R \rangle$.

1.1. Special transformations. For S a nonempty set, we denote by $\mathcal{W}^+(S)$ the free monoid consisting of all words in the alphabet S , and by $\mathcal{W}(S)$ the free monoid of all words in $S \cup S^{-1}$, where S^{-1} denotes a disjoint copy of S containing one letter s^{-1} for each letter s of S . By definition, a group G admits the presentation $\langle S|R \rangle$ if G is isomorphic to $\mathcal{W}(S)/\equiv_R$, where \equiv_R is the least congruence on $\mathcal{W}(S)$ containing the relations of R plus the free group relations $s^{-1}s = ss^{-1} = 1$. For w in $\mathcal{W}(S)$, we denote by \bar{w} the element of $\langle S|R \rangle$ represented by w . We use ε for the empty word, and, for w in $\mathcal{W}(S)$, we denote by w^{-1} the word obtained from w by exchanging the letters s and s^{-1} everywhere and reversing their order. Finally, we say that a word v is a *factor* of another word w if there exist w_1, w_2 satisfying $w = w_1 v w_2$.

Hereafter, we shall consider presentations of a special type.

Definition 1.1. We say that a group presentation (S, R) is *positive* if all relations of R are of the form $v = v'$ —or, equivalently, $v^{-1}v' = 1$ —where v, v' are nonempty words of $\mathcal{W}^+(S)$.

So the point is that the relations can be written with no letter s^{-1} and no empty word. As shows (1), Artin–Tits presentations are positive. If (S, R) is a positive presentation, it makes sense to consider the monoid presented by (S, R) , which will be denoted by $\langle S|R \rangle^+$. In the context of positive presentations, the definition of the group $\langle S|R \rangle$ implies that two words w, w' of $\mathcal{W}(S)$ represent the same element of $\langle S|R \rangle$ if and only if one can go from w to w' by a finite sequence of transformations of the following types:

- (i) Remove some pair $s^{-1}s$ or ss^{-1} , where s is a letter of S ;
- (ii) Replace some factor v by v' , where $v = v'$ is a relation of R ;
- (iii) Insert some pair $s^{-1}s$ or ss^{-1} , where s is a letter of S .

The question we will investigate is the extent to which transformations of the third type (inserting trivial factors $s^{-1}s$ or ss^{-1}) can be avoided. If we consider arbitrary words w, w' , we cannot expect a positive answer, as removing type (iii) transformations yields a relation that is not symmetric. However, the question makes sense when w' is the empty word ε :

Question 1.2. *Assume that (S, R) is a positive presentation and w is a word of $\mathcal{W}(S)$ that represents 1 in $\langle S|R \rangle$. Can one transform w into the empty word by using transformations of type (i) and (ii) only?*

The answer to Question 1.2 is positive when R is empty, that is, in the case of a free group, but the question is not well posed in general. For instance, if $\mathbf{ab} = \mathbf{ba}$ belongs to R , the word $\mathbf{ab}^{-1}\mathbf{a}^{-1}\mathbf{b}$ represents 1, but it is eligible for no transformation of type (i) and (ii). This is due to a lack of symmetry in type (ii) transformations, and it can be fixed by extending the definition so as to also allow replacing v^{-1} by v'^{-1} when $v = v'$ is a relation of R . However, even so, the answer to Question 1.2 remains negative in simple cases. For instance, assume $S = \{\mathbf{a}, \mathbf{b}, \mathbf{c}\}$ and $R = \{\mathbf{ab} = \mathbf{ba}, \mathbf{bc} = \mathbf{cb}, \mathbf{ac} = \mathbf{ca}\}$. Let $w = \mathbf{aBcAbC}$, where we adopt the convention that, in concrete examples, $\mathbf{A}, \mathbf{B}, \dots$ stand for $\mathbf{a}^{-1}, \mathbf{b}^{-1}, \dots$ [16]. Then w represents 1 in $\langle S|R \rangle$, but it is eligible for no transformation of type (i) or (ii).




FIGURE 1. Special transformations viewed in the Cayley graph: relations correspond to oriented cells; special transformations of types 1 and 2 correspond to changing way around a cell: in type 1, the old path (plain) follows one side of the cell whereas the new path (dotted) follows the other side; in type 2_r , the old path crosses the source of the cell, with at least one letter on both sides, and the new path crosses the target instead; type 2_ℓ is symmetric; note that a transformation of type 2_r is not just the inverse of a transformation of type 2_ℓ as the definition requires that the initial words v, v' are not empty whereas it requires nothing for the final words u, u' : for instance, if $v = v'$ is a relation of R , then $v^{-1}v' \overset{2_r}{\rightsquigarrow}_R \varepsilon$ holds, but $\varepsilon \overset{2_\ell}{\rightsquigarrow}_R v^{-1}v'$ does not.

Here comes the point. In addition to the above transformations, we consider a new type of syntactic transformations, namely the *subword reversing* transformations of [11, 12].

Definition 1.3. For (S, R) a positive presentation, we define *special transformations* on words of $\mathcal{W}(S)$ as follows:

- *type 0* (*resp.* ∞): Remove (*resp.* insert) some pair $s^{-1}s$ or ss^{-1} , where s is a letter of S ,
- *type 1*: Replace some factor v by v' , or v^{-1} by v'^{-1} , where $v = v'$ is a relation of R ,
- *type 2_r* (“type 2-right”): Replace some factor $v^{-1}v'$ with uu'^{-1} , where v, v' are nonempty and $vu = v'u'$ is a relation of R ,
- *type 2_ℓ* (“type 2-left”): Replace some factor vv'^{-1} by $u^{-1}u'$, where v, v' are nonempty and $uv = u'v'$ is a relation of R .

For $X \subseteq \{0, 1, 2, \infty\}$, we write $w \overset{X}{\rightsquigarrow}_R w'$ if w can be transformed to w' using special transformations of type belonging to X .

Special transformations are easily illustrated when words are viewed as paths in the Cayley diagram [16], see Figure 1. With the above notation, the standard result is that, for every positive presentation (S, R) and every word w of $\mathcal{W}(S)$, we have

$$(1.1) \quad \overline{w} = 1 \iff w \overset{0,1,\infty}{\rightsquigarrow}_R \varepsilon.$$

Moreover, as is clear from Figure 1, every transformation of type 2 replaces a word with a new word that represents the same element of the involved group, so $w \overset{0,1,2,\infty}{\rightsquigarrow}_R w'$ implies and, therefore, is equivalent to, $\overline{w} = \overline{w'}$. In particular, $w \overset{0,1,2,\infty}{\rightsquigarrow}_R \varepsilon$ is equivalent to $\overline{w} = 1$. The question we will address is whether type ∞ transformations can be avoided:

Question 1.4. Assume that (S, R) is a positive presentation and w is a word of $\mathcal{W}(S)$ that represents 1 in $\langle S|R \rangle$. Does $w \overset{0,1,2}{\rightsquigarrow}_R \varepsilon$ necessarily hold?

Introducing type 2 transformations really changes the situation. For instance, if S is $\{a, b, c\}$ and R consists of $ab = ba$, $bc = cb$, and $ac = ca$, we observed above that, for $w = aBcAbC$, although w represents 1, the relation $w \overset{0,1}{\rightsquigarrow}_R \varepsilon$ fails. Now, $w \overset{0,1,2}{\rightsquigarrow}_R \varepsilon$ holds, since we can write $aBcAbC \overset{2}{\rightsquigarrow}_R aBAbcC \overset{1}{\rightsquigarrow}_R aABcbC \overset{0}{\rightsquigarrow}_R BcbC \overset{1}{\rightsquigarrow}_R BbcC \overset{0}{\rightsquigarrow}_R cC \overset{0}{\rightsquigarrow}_R \varepsilon$.

We are ready to introduce our main definition and state our conjecture.

Definition 1.5. We say that a positive presentation (S, R) satisfies Property H if the relation $w \overset{0,1,2}{\rightsquigarrow}_R \varepsilon$ holds for every word w of $\mathcal{W}(S)$ that represents 1 in $\langle S|R \rangle$.

Conjecture 1.6. *Every Artin–Tits presentation satisfies Property H.*

Before going on, let us immediately note that some positive presentations do *not* satisfy Property H. For instance, let S be $\{a, b, c, d, e, f\}$ and let R consist of the four relations $ca = ace$, $cb = bce$, $da = adf$, and $db = bdf$. Let $w = ACdaBDcb$. Then, as shown in Figure 2, the word w represents 1 in the group $\langle S|R \rangle$, but $w \overset{0,1,2}{\rightsquigarrow}_R \varepsilon$ cannot be true, as w is eligible for no transformation of type 0, 1, or 2. Hence the positive presentation (S, R) does not satisfy Property H.




FIGURE 2. Whenever $ca = ace$, $cb = bce$, $da = adf$, and $db = bdf$ are valid relations, the word $w = ACdaBDcb$ represents 1. However it is eligible for no special transformation, hence cannot satisfy $w \overset{0,1,2}{\rightsquigarrow}_R \varepsilon$.

1.2. Connection with the Dehn algorithm and hyperbolicity. Property H is reminiscent of the property that underlies the Dehn algorithm, and, in this way, it is connected with hyperbolicity. Here, we make this connection precise. The result is stated in Proposition 1.8 below, and it shows that Property H can be viewed as a weak hyperbolicity condition. In this way, Conjecture 1.6 asserts that Artin–Tits groups, which cannot to be hyperbolic in general as they include a free Abelian group of rank two, nevertheless satisfy some weak variant of hyperbolicity.

Hereafter, we denote by $|w|$ the length (number of letters) of a word w . Let us first recall the standard definition of the Dehn algorithm, here in the context of a positive presentation.

Definition 1.7. Assume that (S, R) a positive group presentation. For w, w' in $\mathcal{W}(S)$, we say that w' is obtained from w by a *Dehn transformation with respect to R* if w' is obtained by replacing some factor u of w with a new factor u' such that $|u| > |u'|$ holds and there exists a relation $v = v'$ of R such that $u^{-1}u'$ is a cyclic permutation of $v^{-1}v'$ or of $v'^{-1}v$. We say that $w \overset{0,D}{\rightsquigarrow}_R w'$ holds if w' is obtained from w by a finite sequence of type 0 transformations and Dehn transformations with respect to R .

Clearly, $w \overset{0,D}{\rightsquigarrow}_R w'$ implies $w \equiv_R w'$ and, in particular, $w \overset{0,D}{\rightsquigarrow}_R \varepsilon$ always implies $w \equiv_R \varepsilon$. Saying that the latter implication is an equivalence means that the Dehn algorithm is valid for the presentation (S, R) . It is well known [1, 6, 21] that, in this case, and provided the presentation (S, R) is finite, the group $\langle S|R \rangle$ is hyperbolic in the sense of Gromov.

It turns out that, at least in the case of positive presentations that satisfy simple length constraints, the relations $\overset{0,1,2}{\rightsquigarrow}_R$ and $\overset{0,D}{\rightsquigarrow}_R$ are closely connected.

Proposition 1.8. *Assume that (S, R) is a positive presentation such that, for every relation $v = v'$ of R , we have $||v| - |v'|| \leq 2$. Then, for all w, w' in $\mathcal{W}(S)$, the relation $w \overset{0,D}{\rightsquigarrow}_R w'$ implies $w \overset{0,1,2}{\rightsquigarrow}_R w'$.*

Proof. It is enough to prove that $w \overset{0,1,2}{\rightsquigarrow}_R w'$ holds when w' is obtained from w by one Dehn transformation. As $w \overset{0,1,2}{\rightsquigarrow}_R w'$ implies $w_1 w w_2 \overset{0,1,2}{\rightsquigarrow}_R w_1 w' w_2$ for all w_1, w_2 , it is even sufficient to prove $u \overset{0,1,2}{\rightsquigarrow}_R u'$ when there exists a relation $v = v'$ of R such that $|u| > |u'|$ holds and

$u^{-1}u'$ is a cyclic permutation of $v'^{-1}v$. The case when $u^{-1}u'$ is a cyclic permutation of $v^{-1}v'$ would of course be symmetric.

We consider the possible repartitions of the positive and negative letters in u and u' . If u contains at most one positive and one negative letter, it must contain a factor $s^{-1}t$ or a factor st^{-1} , in which cases it is eligible for a type 2 special transformation, and we directly obtain $u \overset{2}{\rightsquigarrow}_R u'$. Otherwise, assume that u contains only positive letters, that is, it is a factor of v . If u is all of v , then u' is v' , and we obtain $u \overset{1}{\rightsquigarrow}_R u'$. Otherwise, we must have $|u| < |v|$, whence $|u| \leq |v| - 1$, and $|u'| \leq |v| - 2$. By hypothesis, we have $|u| + |u'| = |v| + |v'|$, and we deduce $|v| + |v'| \leq 2|v| - 3$, whence $|v'| \leq |v| - 3$, which is incompatible with the hypothesis on (S, R) . Finally, the case when u contains only negative letters, hence is a factor of v'^{-1} , is symmetric. \square

Thus, at least for presentations satisfying the length constraint of Proposition 1.8, the relation $\overset{0,1,2}{\rightsquigarrow}_R$ is a weak version of $\overset{0,D}{\rightsquigarrow}_R$ and, therefore, Property H is a weak version of hyperbolicity. Artin–Tits presentations are eligible for Proposition 1.8 since all Artin–Tits relations are of the form $v = v'$ with $|v| = |v'|$. As, except in the degenerate case of free groups, Artin–Tits groups are not hyperbolic, Conjecture 1.6 is a weaker statement in the same direction.

Remark 1.9. As it involves the equivalence relation \equiv_R , Property H is connected with the word problem of the considered presentation. However, Property H need *not* imply the decidability of the word problem. Indeed, starting from a word w , there exist in general infinitely many words w' satisfying $w \overset{0,1,2}{\rightsquigarrow}_R w'$. For instance, in the case of 4-strand braids, that is, of the Artin–Tits presentation of type A_3 —which satisfies Property H as will be established in Proposition 1.11 below—there exist infinitely many words w' satisfying $\sigma_2^{-1}\sigma_1^{-1}\sigma_3\sigma_2 \overset{0,1,2}{\rightsquigarrow}_R w'$ [14]. In such a case, there is a priori no method for establishing the failure of $w \overset{0,1,2}{\rightsquigarrow}_R \varepsilon$, and therefore, when Property H is satisfied, for establishing the failure of $w \equiv_R \varepsilon$.

To complete the discussion of the connection between Property H and the Dehn algorithm, we now briefly mention the approach of Appel and Schupp [3] in extra-large type, and explain why it does not seem to naturally lead to a proof of Property H.

An Artin–Tits presentation (S, R) is said to be of *extra-large type* if all relations of R have the form $v = v'$ with $|v| = |v'| \geq 4$. Appel and Schupp [3] prove that, if (S, R) is an Artin–Tits presentation of extra-large type and, if \widehat{R} is the set of all relations $u = u'$ where u, u' are R -equivalent words of $\mathcal{W}(S)$ involving only two letters of S and such that $u^{-1}u'$ is cyclically reduced, then a word w of $\mathcal{W}(S)$ represents 1 in $\langle S|R \rangle$ if and only if $w \overset{0,D}{\rightsquigarrow}_{\widehat{R}} \varepsilon$ holds (here we do not assume that the words u, u' are positive, so the presentation (S, \widehat{R}) is not positive in general). In other words, the Dehn algorithm with respect to the family \widehat{R} solves the word problem, a result that does not contradict the non-hyperbolicity of the group $\langle S|R \rangle$ as \widehat{R} is always infinite. Owing to Proposition 1.8, it is natural to wonder whether the implication

$$(1.2) \quad w \overset{0,D}{\rightsquigarrow}_{\widehat{R}} w' \quad \Rightarrow \quad w \overset{0,1,2}{\rightsquigarrow}_R w'$$

is always true. A direct proof of this implication would provide a proof of Conjecture 1.6, at least in the extra-large type: if w represents 1, then $w \overset{0,D}{\rightsquigarrow}_{\widehat{R}} \varepsilon$ holds, and we would deduce $w \overset{0,1,2}{\rightsquigarrow}_R \varepsilon$. However, (1.2) need not be true, even in very simple cases, as the following counter-example shows.

Example 1.10. Consider $S = \{a, b\}$ and let R consist of the unique relation $abab = baba$. Then (S, R) is an Artin–Tits presentation of extra-large type, and the associated Coxeter

group is the eight element dihedral group. Let $w = \mathbf{Ab}^2\mathbf{abaB}$ and $w' = \mathbf{ababA}$. Then w and w' are R -equivalent, we have $|w| = 7 > |w'| = 5$. So $w \overset{0,D}{\rightsquigarrow}_R w'$ holds. We claim that $w \overset{0,1,2}{\rightsquigarrow}_R w'$ fails. The argument is as follows. We consider the finite fragment Γ of the Cayley graph of $\langle S|R \rangle$ represented in plain lines in Figure 3: the vertices are the nine left-divisors of the element \mathbf{abab}^2 in the monoid $\langle S|R \rangle^+$. Then the word w is, in the sense of [10] or [13, Chapter IV], traced from \mathbf{a} in Γ , that is, there exists inside Γ a path starting from the vertex labeled \mathbf{a} whose edges wear as labels the successive letters of w . As explained in the cited references, the family of all words traced from a given vertex in the fragment of the Cayley graph made of all left-divisors of a given element is closed under special transformations of types 0, 1, and 2. Now the picture shows that w' is *not* traced from \mathbf{a} in Γ and, therefore, $w \overset{0,1,2}{\rightsquigarrow}_R w'$ cannot hold.




FIGURE 3. Proving that $w \rightsquigarrow w'$ fails in Example 1.10: the word w is traced from the vertex \mathbf{a} in the fragment of the Cayley graph written in plain lines, and so is every word obtained from w using special transformations, whereas w' is not traced from the vertex \mathbf{a} in that fragment.

1.3. The case of spherical type. Artin–Tits presentations of spherical type (also known as of finite type), that is, those such that the associated Coxeter group is finite [18], provide natural test-cases: the corresponding groups are well understood and any conjecture involving general Artin–Tits groups can be expected to be provable for spherical type. This is the case for Property H.

Proposition 1.11. *Every Artin–Tits presentation of spherical type satisfies Property H.*

In order to establish Proposition 1.11 and to prepare the arguments of Subsection 2.2, we recall some terminology about divisibility relations. If M is a monoid and f, g belong to M , one says that f is a *left-divisor* of g , and that g is a *right-multiple* of f , if $fg' = g$ holds for some element g' of M . If M has no nontrivial (that is, distinct of 1) invertible element—as is the case of every monoid $\langle S|R \rangle^+$ defined by a positive presentation—the left-divisibility relation is a partial ordering on M . Right-divisors are defined symmetrically: f is a *right-divisor* of g , and g is a *left-multiple* of f , if $g = g'f$ holds for some g' . Finally, two elements of M are called *left-coprime* (*resp.* *right-coprime*) if they admit no nontrivial common left-divisor (*resp.* right-divisor).

The specificity of the spherical type is the following classical result describing $\langle S|R \rangle$ as a group of (right)-fractions.

Proposition 1.12. [5, 16, 23, 7] *Assume that (S, R) is an Artin–Tits presentation of spherical type. Then the monoid $\langle S|R \rangle^+$ embeds in the group $\langle S|R \rangle$ and, for every element g of $\langle S|R \rangle$, there exists a unique pair (g_1, g_2) of right-coprime elements of $\langle S|R \rangle^+$ satisfying $g = g_1g_2^{-1}$. Moreover, if we have $g = f_1f_2^{-1}$ with f_1, f_2 in $\langle S|R \rangle^+$, there exists h in $\langle S|R \rangle^+$ such that $f_e = g_e h$ holds for $e = 1, 2$.*

In the above context, the elements g_1 and g_2 above are called the *right-numerator* and the *right-denominator* of g , here denoted by $N_r(g)$ and $D_r(g)$. In terms of words, there exists a simple connection with special transformations.

Lemma 1.13. [9] *Assume that (S, R) is a Artin-Tits presentation of spherical type and w is a word of $\mathcal{W}(S)$. Let g_1, g_2 be the right numerator and denominator of \bar{w} .*

- (i) *There exist w_1, w_2 in $\mathcal{W}^+(S)$ satisfying $w \overset{0,2}{\rightsquigarrow}_R w_1 w_2^{-1}$ and $\bar{w}_e = g_e$ for $e = 1, 2$.*
- (ii) *For all w'_1, w'_2 in $\mathcal{W}^+(S)$ satisfying $\bar{w}'_e = g_e$ for $e = 1, 2$, we have $w \overset{0,1,2}{\rightsquigarrow}_R w'_1 w'^{-1}_2$.*

Proof (sketch). (i) It follows from the fact that any two elements in the monoid $\langle S|R \rangle^+$ admit a common left-multiple that, starting from w and performing type 0 and 2_ℓ special transformations, one arrives in finitely many steps to a word of the form $v_1^{-1} v_2$ with v_1, v_2 in $\mathcal{W}^+(S)$. By a symmetric argument, starting from $v_1^{-1} v_2$ and performing type 0 and 2_r special transformations, one arrives in finitely many steps to a word of the form $w_1 w_2^{-1}$ with w_1, w_2 in $\mathcal{W}^+(S)$. So $w \overset{0,2}{\rightsquigarrow}_R w_1 w_2^{-1}$ holds. Moreover, the element of $\langle S|R \rangle^+$ represented by $v_1 w_1$ and $v_2 w_2$ is the least common right-multiple of \bar{v}_1 and \bar{v}_2 , which implies that \bar{w}_1 and \bar{w}_2 are right-coprime. The uniqueness result in Proposition 1.12 then implies that w_e represents g_e for $e = 1, 2$.

(ii) Let w'_e be any word of $\mathcal{W}^+(S)$ representing g_e . The uniqueness result of Proposition 1.12 implies that $w'_e \equiv_R^+ w_e$, where w_1, w_2 are the words obtained in (i). Then we have $w_1 w_2^{-1} \overset{1}{\rightsquigarrow}_R w'_1 w'^{-1}_2$. By composing, we deduce $w \overset{0,1,2}{\rightsquigarrow}_R w'_1 w'^{-1}_2$. \square

Due to the symmetry of Artin-Tits relations, Proposition 1.12 and Lemma 1.13 have left counterparts involving left-fractions of the form $g_1^{-1} g_2$ and negative-positive words $w_1^{-1} w_2$. We shall then speak of the left-numerator and the left-denominator of g , denoted by $D_\ell(g)$ and $N_\ell(g)$. The latter are left-coprime elements of $\langle S|R \rangle^+$.

We can now easily complete the argument.

Proof of Proposition 1.11. Assume that w is a word of $\mathcal{W}(S)$ that represents 1 in $\langle S|R \rangle$. Then the left numerator and denominator of \bar{w} are 1. By Lemma 1.13, there exist w_1, w_2 in $\mathcal{W}^+(S)$, both representing 1 in $\langle S|R \rangle^+$, and satisfying $w \overset{0,2}{\rightsquigarrow}_R w_1 w_2^{-1}$. The only possibility for a word of $\mathcal{W}^+(S)$ to represent 1 is to be empty. So we must have $w \overset{0,2}{\rightsquigarrow}_R \varepsilon$ and, a fortiori, $w \overset{0,1,2}{\rightsquigarrow}_R \varepsilon$. So Property H is satisfied. \square

Remark 1.14. The result that $w \equiv_R \varepsilon$ implies $w \overset{0,2}{\rightsquigarrow}_R \varepsilon$ is false for general Artin-Tits presentations. For instance, consider the (right-angled) presentation (S, R) with $S = \{a, b, c, d\}$ and $R = \{ac = ca, bc = cb, ad = da, bd = db\}$. The associated group is a direct product of two free groups of rank 2, respectively generated by a, b and c, d . Let $w = ACdaBDcb$. Then $w \overset{0,1,2}{\rightsquigarrow}_R \varepsilon$ is true, as we can for instance write

$$\begin{aligned} ACdaBDcb &\overset{1}{\rightsquigarrow}_R CAdaBDcb \overset{1}{\rightsquigarrow}_R CAadBDcb \\ &\overset{0}{\rightsquigarrow} CdBDcb \overset{1}{\rightsquigarrow}_R CdDBcb \overset{0}{\rightsquigarrow} CBcb \overset{1}{\rightsquigarrow}_R CBbc \overset{0}{\rightsquigarrow} Cc \overset{0}{\rightsquigarrow} \varepsilon, \end{aligned}$$

hence w represents 1 in $\langle S|R \rangle$. On the other hand, $w \overset{0,2}{\rightsquigarrow}_R \varepsilon$ is false, as w is eligible for no transformation of type 0 or 2.

1.4. An application. We conclude the section with an application of Property H to the embeddability of the monoid $\langle S|R \rangle^+$ in the group $\langle S|R \rangle$. This application provides an additional motivation for investigating Property H, especially in the case of Artin-Tits presentations. As the proof appears in [12] (in a slightly different setting), we shall only state the result here.

To this end, we first need to introduce a condition called completeness. We recall that $\mathcal{W}^+(S)$ denotes the family of all words in $\mathcal{W}(S)$ that contain no letter s^{-1} . Then, by definition, the monoid $\langle S|R \rangle^+$ (isomorphic to) $\mathcal{W}^+(S)/\equiv_R^+$, where \equiv_R^+ is the least congruence on $\mathcal{W}^+(S)$ that includes R . For w, w' in $\mathcal{W}^+(S)$, by definition, $w \equiv_R^+ w'$ is equivalent to $w \overset{1}{\rightsquigarrow}_R w'$ and, from there, to $w^{-1}w' \overset{0,1}{\rightsquigarrow}_R \varepsilon$. Completeness is a formally similar condition that involves special transformations of type 2_r instead of type 1.

Definition 1.15. [11, 12] Assume that (S, R) is a positive presentation. We say that (S, R) is *complete with respect to right-reversing* if, for all w, w' in $\mathcal{W}^+(S)$, the relation $w \equiv_R^+ w'$ is equivalent to $w^{-1}w' \overset{0,2_r}{\rightsquigarrow}_R \varepsilon$.

That $w^{-1}w' \overset{0,2_r}{\rightsquigarrow}_R \varepsilon$ implies $w \equiv_R^+ w'$ is always true, so the non-trivial part is the converse implication, namely the condition that types 0 and 2_r transformations are sufficient to detect all pairs of \equiv_R^+ -equivalent words. The embeddability result we wish to mention is as follows.

Proposition 1.16. [12, Proposition 3.29] *Assume that (S, R) is a positive presentation that is complete with respect to right-reversing, contains no relation of the form $sv = sv'$ with $v \neq v'$, and satisfies Property H. Then the monoid $\langle S|R \rangle^+$ embeds in the group $\langle S|R \rangle$.*

The (non-trivial) proof consists in showing that, if two words w, w' of $\mathcal{W}^+(S)$ satisfy $w \equiv_R^+ w'$, then one can go from w to w' using a certain “fan” of words of $\mathcal{W}^+(S)$ that are connected under \equiv_R^+ . One then deduces that $w \equiv_R^+ w'$ implies $w \equiv_R^+ w'$, which means that $\langle S|R \rangle^+$ embeds in $\langle S|R \rangle$.

Note that, if (S, R) is eligible for Proposition 1.16, then the monoid $\langle S|R \rangle^+$ must be left- and right-cancellative, that is, $fg = fg'$ and $gh = g'h$ both imply $g = g'$. It is easy to see that completeness plus the hypothesis on relations $sv = sv'$ implies left-cancellativity. But these conditions alone say nothing about right-cancellativity. This shows that, in such a context, Property H includes a hidden right-cancellativity condition.

For our current purpose, the point is that every Artin–Tits presentation is complete with respect to right-reversing [11], a result that in essence goes back to Garside [17] in the case of classical braids, that is, for Artin–Tits presentations of type A, and to Brieskorn and Saito [5] in the general case. It follows that, if Artin–Tits presentations satisfy Property H, they are eligible for Proposition 1.16. L. Paris [22] proved that, if (S, R) is an Artin–Tits presentation, then $\langle S|R \rangle^+$ always embeds in $\langle S|R \rangle$. The beautiful but indirect argument of [22] uses certain linear representations extending the Lawrence–Krammer representation of braids [20], and a more direct proof remains to be discovered. Together with Proposition 1.16, a proof of Conjecture 1.6 would arguably provide such a natural proof.

2. ARTIN–TITS GROUPS OF TYPE FC

The family of Artin–Tits groups of type FC is the smallest class of groups that contains all Artin–Tits groups of spherical type and is closed under forming an amalgamated product over a standard parabolic subgroup. The aim of this section is to prove

Proposition 2.1. *Every Artin–Tits presentation of type FC satisfies Property H.*

This result shows that Property H is satisfied for a large number of Artin–Tits presentations.

The section is organized as follows. In Subsection 2.1, we introduce a strengthening H^\sharp of Property H that is more suitable for amalgamated products. Then we show that all Artin–Tits presentations of spherical type satisfy Property H^\sharp in Subsection 2.2. Finally, we develop in Subsection 2.3 the inductive argument needed to establish Proposition 2.1.

The general scheme and the techniques used in this section follow those of Altobelli [2]. However, the argument has to be revisited completely because we are considering special transformations and the relation $\overset{0,1,2}{\rightsquigarrow}_R$ instead of \equiv_R . Also, in Subsection 2.3, we do not restrict to Artin–Tits groups and propose a new, hopefully optimized construction.

2.1. Property H^\sharp . The definition of the family of Artin–Tits presentations of type FC as the closure of the family of Artin–Tits presentations of spherical type under a certain type of amalgamated product naturally suggests an inductive proof for Proposition 2.1. However, we first need to modify Property H to make the induction possible.

By definition of standard parabolic subgroups [24], the amalgamated products involved in the construction of type FC groups correspond, in terms of presentations, to a union.

Lemma 2.2. *Say that two presentations (S_1, R_1) and (S_2, R_2) are in amalgam position if, denoting by G'_e the subgroup of $\langle S_e | R_e \rangle$ generated by $S_1 \cap S_2$, the identity map of $S_1 \cap S_2$ induces an isomorphism of G'_1 onto G'_2 . Then $(S_1 \cup S_2, R_1 \cup R_2)$ is a presentation of the amalgamated product $\langle S_1 | R_1 \rangle *_{G'_1=G'_2} \langle S_2 | R_2 \rangle$.*

It directly follows from the results of van der Lek [24] that two Artin–Tits presentations (S_1, R_1) , (S_2, R_2) are in amalgam position if and only if the relations of R_1 and of R_2 that involve generators in $S_1 \cap S_2$ coincide. Thus, owing to Proposition 1.11, which says that all Artin–Tits presentations of spherical type satisfy Property H, the natural approach for proving Proposition 2.1 should be to establish that the union of two Artin–Tits presentations that are in amalgam position and satisfy Property H again satisfies Property H. This however seems difficult, even probably false in general. What we shall do below is to introduce some strengthening H^\sharp of H and apply the expected induction scheme to Property H^\sharp rather than to Property H.

Hereafter, if (S, R) is a positive presentation and S' is a subset of S , we denote by $\langle\langle S' \rangle\rangle$ the subgroup of $\langle S | R \rangle$ generated by S' . Similarly, we denote by $\langle\langle S' \rangle\rangle^+$ the submonoid of $\langle S | R \rangle^+$ generated by S' . If G' is a subgroup of $\langle S | R \rangle$, a *(left)- G' -transversal* is a subset of $\langle S | R \rangle$ that contains one element exactly in each left- G' -coset, and that contains 1—the latter condition is not always required, but here we explicitly include it in our definition. Below, we shall consider families of transversals with respect to various subgroups: we define an *S -sequence of transversals* in $\langle S | R \rangle$ to be a sequence $\mathbf{T} = (\mathbf{T}(S'))_{S' \subseteq S}$ such that $\mathbf{T}(S')$ is $\langle\langle S' \rangle\rangle$ -transversal in $\langle S | R \rangle$ for every S' included in S .

Definition 2.3. Assume that (S, R) is a positive presentation and \mathbf{T} is an S -sequence of transversals in $\langle S | R \rangle$. We say that (S, R) satisfies *Property H^\sharp with respect to \mathbf{T}* if, for all S', S_0 included in S and w in $\mathcal{W}(S)$, the following relation $\mathcal{P}_{\mathbf{T}}(S', S_0, w)$ is satisfied:

$$(2.1) \quad \overline{w} \in \langle\langle S' \rangle\rangle \text{ implies } \exists v \in \mathcal{W}(S') \exists u \in \mathcal{W}(S' \cap S_0) (w \overset{0,1,2}{\rightsquigarrow}_R vu \text{ and } \overline{v} \in \mathbf{T}(S_0)).$$

Roughly speaking, $\mathcal{P}_{\mathbf{T}}(S', S_0, w)$ says that, starting with a word w and using special transformations, we can isolate a representative of the (unique) element of the distinguished transversal $\mathbf{T}(S_0)$ that lies in the left- $\langle\langle S_0 \rangle\rangle$ -coset of \overline{w} , and, if \overline{w} lies in $\langle\langle S' \rangle\rangle$, we can moreover require that the final words lie in $\mathcal{W}(S')$.

Lemma 2.4. *Property H^\sharp implies Property H.*

Proof. Assume that (S, R) satisfies Property H^\sharp with respect to \mathbf{T} , and that w represents 1 in $\langle S | R \rangle$. Then \overline{w} belongs to $\langle\langle \emptyset \rangle\rangle$. Hence $\mathcal{P}_{\mathbf{T}}(\emptyset, S, w)$ says that there exists v and u in $\mathcal{W}(\emptyset)$ such that $w \overset{0,1,2}{\rightsquigarrow}_R vu$ holds. Now the only element of $\mathcal{W}(\emptyset)$ is the empty word ε , so the only possibility is $v = u = \varepsilon$, whence $w \overset{0,1,2}{\rightsquigarrow}_R \varepsilon$. So (S, R) satisfies Property H. \square

The result we shall establish is then

Proposition 2.5. *Every Artin–Tits presentation of type FC satisfies Property H^\sharp .*

By Lemma 2.4, Proposition 2.5 directly implies Proposition 2.1. The proof of Proposition 2.5 will occupy the rest of this section, and consists in proving that every Artin–Tits presentation of spherical type satisfies Property H^\sharp (Proposition 2.7) and that the union of two presentations that are in amalgam position and satisfy Property H^\sharp satisfies Property H^\sharp (Proposition 2.14). We begin with two direct consequences of the definition:

Lemma 2.6. *Assume that (S, R) is a positive presentation that satisfies Property H^\sharp .*

- (i) *For all S', S'' included in S , we have $\langle\langle S' \rangle\rangle \cap \langle\langle S'' \rangle\rangle = \langle\langle S' \cap S'' \rangle\rangle$.*
- (ii) *For all S' included in S and w in $\mathcal{W}(S)$,*

$$(2.2) \quad \bar{w} \in \langle\langle S' \rangle\rangle \quad \text{implies} \quad \exists v \in \mathcal{W}(S') \quad (w \overset{0,1,2}{\rightsquigarrow}_R v).$$

Proof. Assume that (S, R) satisfies Property H^\sharp with respect to \mathbf{T} .

(i) The inclusion $\langle\langle S' \cap S'' \rangle\rangle \subseteq \langle\langle S' \rangle\rangle \cap \langle\langle S'' \rangle\rangle$ is always true. For the other inclusion, assume that g belongs to $\langle\langle S' \rangle\rangle \cap \langle\langle S'' \rangle\rangle$. Let w be a word of $\mathcal{W}(S)$ representing g . By $\mathcal{P}_{\mathbf{T}}(S', S'', w)$, there exist v in $\mathcal{W}(S')$ and u in $\mathcal{W}(S' \cap S'')$ satisfying $w \overset{0,1,2}{\rightsquigarrow}_R vu$ and $\bar{v} \in \mathbf{T}(S'')$. Then we have $g = \bar{w} = \bar{v}\bar{u}$. By hypothesis \bar{v} belongs to $\langle\langle S' \rangle\rangle$, so \bar{v} is the element of the $\langle\langle S'' \rangle\rangle$ -transversal $\mathbf{T}(S'')$ that belongs to the left- $\langle\langle S'' \rangle\rangle$ -coset of \bar{w} . Now, by hypothesis, \bar{w} belongs to $\langle\langle S'' \rangle\rangle$, hence \bar{v} must be 1, and, therefore, we have $\bar{w} = \bar{u}$, whence $g = \bar{u} \in \langle\langle S' \cap S'' \rangle\rangle$.

(ii) Assume $\bar{w} \in \langle\langle S' \rangle\rangle$. Then, by $\mathcal{P}_{\mathbf{T}}(S', \emptyset, w)$, there exist v in $\mathcal{W}(S')$ and u in $\mathcal{W}(\emptyset)$ satisfying $w \overset{0,1,2}{\rightsquigarrow}_R vu$ and $\bar{v} \in \mathbf{T}(\emptyset)$. The only possibility is $u = \varepsilon$, and, therefore, we have $w \overset{0,1,2}{\rightsquigarrow}_R v$. So (2.2) is satisfied. \square

2.2. The spherical case. According to the scheme described above, the first step toward Proposition 2.5 is to consider Artin–Tits presentations of spherical type. We shall establish

Proposition 2.7. *Every Artin–Tits presentation of spherical type satisfies Property H^\sharp .*

A first observation is that, for Artin–Tits presentations, Condition $\mathcal{P}_{\mathbf{T}}(S', S_0, w)$, which simultaneously involves two subsets S_0, S' of S , can be split into two simpler conditions involving only one subset of S at a time.

Lemma 2.8. *Assume that (S, R) is an Artin–Tits presentation. Then (S, R) satisfies Property H^\sharp with respect to \mathbf{T} if and only if, for all S_0 included in S and w in $\mathcal{W}(S)$, we have (2.2) and*

$$(2.3) \quad \exists v \in \mathcal{W}(S) \exists u \in \mathcal{W}(S_0) \quad (w \overset{0,1,2}{\rightsquigarrow}_R vu \text{ and } \bar{v} \in \mathbf{T}(S_0)).$$

Proof. We already observed in Lemma 2.6 that Property H^\sharp implies (2.2) for all S' and w . On the other hand, (2.3) is (equivalent to) $\mathcal{P}_{\mathbf{T}}(S, S_0, w)$, so Property H^\sharp implies (2.2) and (2.3).

Conversely, assume that (S, R) is an Artin–Tits presentation that satisfies (2.2) and (2.3) for all S', S_0 , and w . Let S_0, S' be subsets of S and let w be a word of $\mathcal{W}(S)$ that represents an element of $\langle\langle S' \rangle\rangle$. By (2.2), there exists a word w' of $\mathcal{W}(S')$ satisfying $w \overset{0,1,2}{\rightsquigarrow}_R w'$ holds. Next, by (2.3), we find v in $\mathcal{W}(S)$ and u in $\mathcal{W}(S_0)$ satisfying $w' \overset{0,1,2}{\rightsquigarrow}_R vu$ and $\bar{v} \in \mathbf{T}(S_0)$. Now, due to the specific syntactic form of Artin–Tits relations, the hypothesis that w' lies in $\mathcal{W}(S')$ implies that w'' lies in $\mathcal{W}(S')$ for every word w'' satisfying $w' \overset{0,1,2}{\rightsquigarrow}_R w''$. Indeed, a special transformation of type 0, 1, or 2 can never let a new letter of S or S^{-1} appear. Hence v must lie in $\mathcal{W}(S')$, and u must lie in $\mathcal{W}(S_0 \cap S')$. So $\mathcal{P}_{\mathbf{T}}(S', S_0, w)$ is satisfied. \square

Next, we observe that (2.2) is almost trivial.

Lemma 2.9. *Assume that (S, R) is an Artin–Tits presentation of spherical type. Then (2.2) is satisfied for all S_0 included in S and w in $\mathcal{W}(S)$.*

Proof. Assume that S_0 is included in S and w is an element of $\mathcal{W}(S)$ that represents an element of $\langle\langle S_0 \rangle\rangle$. Then \bar{w} has at least one representative that lies in $\mathcal{W}(S_0)$ and, therefore, $\mathfrak{N}_r(\bar{w})$ and $\mathfrak{D}_r(\bar{w})$ admit representatives, say v_1, v_2 that lie in $\mathcal{W}^+(S_0)$. Then the word $v_1 v_2^{-1}$ is a word of $\mathcal{W}(S_0)$ and, by Lemma 1.13(ii), we have $w \overset{0,1,2}{\rightsquigarrow}_R v_1 v_2^{-1}$. \square

So we are left with proving that, if (S, R) is an Artin-Tits presentation of spherical type, then all instances of (2.3) are satisfied. We begin with a general auxiliary result that will be used several times in the sequel.

Lemma 2.10. *Assume that (S, R) is an Artin-Tits presentation, S_0 is included in S , and we have $gh = g'h'$ with g, g' in $\langle S|R \rangle^+$ and h, h' in $\langle\langle S_0 \rangle\rangle^+$. If g is right-divisible by no nontrivial element of $\langle\langle S_0 \rangle\rangle^+$, then necessarily h right-divides h' .*

Proof. (See Figure 4.) In the monoid $\langle S|R \rangle^+$, the elements h and h' admit a least common left-multiple, say h'' , that lies in $\langle\langle S_0 \rangle\rangle^+$. Since gh , which is also $g'h'$, is a left-multiple of h and h' , it is a left-multiple of h'' , say $gh = g''h''$ with $g'' \in \langle S|R \rangle^+$. Write $h'' = fh$. We find $gh = g''fh$, whence $g = g''f$. By construction, f belongs to $\langle\langle S_0 \rangle\rangle^+$ and right-divides g , so the hypothesis that g is right-divisible by no nontrivial element of $\langle\langle S_0 \rangle\rangle^+$ implies $f = 1$, whence $h'' = h$. The latter equality means that h' right-divides h . \square




FIGURE 4. Proof of Lemma 2.10.

Until the end of this subsection, we assume that (S, R) is an Artin-Tits presentation of spherical type. We denote by G the group $\langle S|R \rangle$, and by G^+ the monoid $\langle S|R \rangle^+$. We shall now construct an S -sequence of transversals \mathbf{T} with the aim of proving that (2.3) is always satisfied.

Definition 2.11. Assume that S_0 is included in S . An element g of G is called S_0 -minimal if, for every h in $\langle\langle S_0 \rangle\rangle$, we have $|\mathfrak{D}_\ell(gh)| \geq |\mathfrak{D}_\ell(g)|$ and $|\mathfrak{D}_\ell(gh)| = |\mathfrak{D}_\ell(g)|$ implies $|\mathfrak{N}_\ell(gh)| \geq |\mathfrak{N}_\ell(g)|$. Then we put

$$\mathbf{T}(S_0) = \{g \in G \mid g \text{ is } S_0\text{-minimal}\}.$$

We shall show that \mathbf{T} is an S -sequence of transversals witnessing for Property H[#]. We begin with several technical results about S_0 -minimal elements.

Lemma 2.12. (i) *An element of G^+ is S_0 -minimal if and only if it is right-divisible by no nontrivial element of $\langle\langle S_0 \rangle\rangle^+$.*

(ii) *If g is S_0 -minimal, then, for every h in $\langle\langle S_0 \rangle\rangle^+$, the elements $\mathfrak{D}_\ell(g)$ and $\mathfrak{N}_\ell(g)h$ are left-coprime.*

(iii) *If g is S_0 -minimal and lies in G^+ , then, for every h in $\langle\langle S_0 \rangle\rangle$, the elements $\mathfrak{N}_r(gh)$ and $\mathfrak{D}_r(h)$ are right-coprime.*

(iv) *If g is S_0 -minimal, then so is $\mathfrak{N}_\ell(g)$.*

Proof. (i) Assume that g is an element of G^+ and h is an element of $\langle\langle S_0 \rangle\rangle^+$ that right-divides g , say $g = g'h$ with $g' \in G^+$. Then we have $|\mathfrak{D}_\ell(gh^{-1})| = |\mathfrak{D}_\ell(g)| = 0$ and $|\mathfrak{N}_\ell(gh^{-1})| = |g'| = |g| - |h|$. If g is S_0 -minimal, the only possibility is $|h| = 0$, that is, $h = 1$.

Conversely, assume that g is an element of G^+ that is right-divisible by no nontrivial element of $\langle\langle S_0 \rangle\rangle^+$. Let h belong to $\langle\langle S_0 \rangle\rangle$. We have $\text{D}_\ell(g) = 1$, so $|\text{D}_\ell(gh)| \geq |\text{D}_\ell(g)|$ is always satisfied. Assume $|\text{D}_\ell(gh)| = |\text{D}_\ell(g)|$, hence $|\text{D}_\ell(gh)| = 0$, that is, $gh \in G^+$. Let $g' = gh$, and write $h = h_1 h_2^{-1}$ with h_1, h_2 in $\langle\langle S_0 \rangle\rangle^+$. Then we have $gh_1 = g'h_2$. By Lemma 2.10, h_2 right-divides h_1 , that is, h belongs to $\langle\langle S_0 \rangle\rangle^+$. Then we obtain $|\text{N}_\ell(gh)| = |gh| \geq |g| = |\text{N}_\ell(g)|$, and g is S_0 -minimal.

(ii) Assume that g is S_0 -minimal and h lies in $\langle\langle S_0 \rangle\rangle^+$. Put $g_1 = \text{D}_\ell(g)$, $g_2 = \text{N}_\ell(g)$. By definition of an S_0 -minimal element, we have $|\text{D}_\ell(gh)| \geq |\text{D}_\ell(g)| = |g_1|$. Assume that f is a common left-divisor of g_1 and $g_2 h$, say $g_1 = f g'_1$ and $g_2 h = f g'_2$ with g'_1, g'_2 in G^+ . As we have $gh = g_1^{-1} g'_2$, we deduce $|\text{D}_\ell(gh)| \leq |g'_1| = |g_1| - |f|$. The only possibility is $|f| = 0$, that is, $f = 1$. Hence g_1 and $g_2 h$ are left-coprime.

(iii) Let $h_1 = \text{N}_r(h)$ and $h_2 = \text{D}_r(h)$. Assume that f right-divides gh_1 and h_2 , say $gh_1 = g'f$. As f right-divides h_2 , it lies in $\langle\langle S_0 \rangle\rangle^+$. As g is S_0 -minimal, (i) implies that it is right-divisible by no nontrivial element of $\langle\langle S_0 \rangle\rangle^+$. Then Lemma 2.10 implies that f right-divides h_1 and, therefore, the hypothesis that h_1 and h_2 are right-coprime implies $f = 1$. So gh_1 and h_2 are right-coprime, as expected.

(iv) Assume that g is S_0 -minimal. Let $g_1 = \text{D}_\ell(g)$ and $g_2 = \text{N}_\ell(g)$. Assume $g_2 = g'h$ with h in $\langle\langle S_0 \rangle\rangle^+$. Let $g' = g_1^{-1} g' = gh^{-1}$. The hypothesis that g_1 and g_2 are left-coprime implies that g_1 and g' are left-coprime. Hence we have $\text{D}_\ell(g') = g_1 = \text{D}_\ell(g)$ and $\text{N}_\ell(g') = g'$. As we have $g' = gh^{-1}$ with $h^{-1} \in \langle\langle S_0 \rangle\rangle$ and $|\text{D}_\ell(g')| = |\text{D}_\ell(g)|$, the hypothesis that g is S_0 minimal implies $|\text{N}_\ell(g')| \geq |\text{N}_\ell(g)|$, that is, $|g'| \geq |g_2|$. As we have $|g_2| = |g'| + |h|$, the only possibility is $|h| = 1$, that is, $h = 1$. Thus g has no nontrivial right-divisor in $\langle\langle S_0 \rangle\rangle$. By (i), g is S_0 -minimal. \square

Lemma 2.13. *For every S_0 included in S , the family $\mathbf{T}(S_0)$ is a $\langle\langle S_0 \rangle\rangle$ -transversal in G .*

Proof. Let C be a left- $\langle\langle S_0 \rangle\rangle$ -coset. Let C' be the set $\{g \in C \mid |\text{D}_\ell(g)| \text{ is minimal}\}$, and C'' be $\{g \in C' \mid |\text{N}_\ell(g)| \text{ is minimal}\}$. By definition, C'' is nonempty, and every element of C'' is an element of C that is $\langle\langle S_0 \rangle\rangle$ -minimal. So every left- $\langle\langle S_0 \rangle\rangle$ -coset contains at least one S_0 -minimal element.

Assume now that g and g' belong to $\mathbf{T}(S_0)$ and lie in the same left- $\langle\langle S_0 \rangle\rangle$ -coset, say $g = g'h$ with $h \in \langle\langle S_0 \rangle\rangle$. Let g_1, g_2 be the left denominator and numerator of g , and g'_1, g'_2 be those of g' . Write $h = h_1 h_2^{-1}$ with h_1, h_2 in $\langle\langle S_0 \rangle\rangle^+$. As g' is S_0 -minimal, Lemma 2.12(ii) implies that g'_1 and $g'_2 h_2$ are left-coprime. Now we have $g_1^{-1} (g_2 h_1) = g_1^{-1} (g'_1 h_2)$. As g'_1 and $g'_2 h_2$ are left-coprime, g'_1 must right-divide g_1 . By a symmetric argument exchanging the roles of g and g' and those of h_1 and h_2 , we see that g_1 must right-divide g'_1 . We deduce $g_1 = g'_1$, whence $g_2 h_1 = g'_2 h_2$. By Lemma 2.12(iv), g_2 is S_0 -minimal, so, by Lemma 2.12(i), g_2 is right-divisible by no nontrivial element of $\langle\langle S_0 \rangle\rangle^+$. Hence, by Lemma 2.10, h_2 right-divides h_1 . A symmetric argument shows that h_1 right-divides h_2 . We deduce $h_1 = h_2$, whence $g = g'$. So every left- $\langle\langle S_0 \rangle\rangle$ -coset contains at most one S_0 -minimal element.

Finally, it is clear that 1 is S_0 -minimal, hence $\mathbf{T}(S_0)$ contains 1. \square

With these results at hand, we can complete the argument.

Proof of Proposition 2.7. We have seen that the only point that remains to be proved is the satisfaction of (2.3) for all S_0 and w . So, assume that S_0 are included in S and w is a word of $\mathcal{W}(S)$. Put $g = \overline{w}$. As $\mathbf{T}(S_0)$ is $\langle\langle S_0 \rangle\rangle$ -transversal, we have $g = g'h$ for some (unique) g' in $\mathbf{T}(S_0)$ and h in $\langle\langle S_0 \rangle\rangle$. In G , we have

$$(2.4) \quad \text{D}_\ell(g)^{-1} \text{N}_\ell(g) \text{D}_r(h) = g \text{D}_r(h) = g' h \text{D}_r(h) = \text{D}_\ell(g')^{-1} \text{N}_\ell(g') \text{N}_r(h).$$

By hypothesis, g' is S_0 -minimal. Hence, by Lemma 2.12(ii), $\text{D}_\ell(g')$ and $\text{N}_\ell(g') \text{N}_r(h)$ are left-coprime. So, by (the left counterpart of) Proposition 1.12, (2.4) implies the existence of f

in G^+ satisfying

$$D_\ell(g) = fD_\ell(g') \quad \text{and} \quad N_\ell(g)D_r(h) = fN_\ell(g')N_r(h),$$

see Figure 5. By Lemma 2.12(iv), the hypothesis that g' is S_0 -minimal implies that $N_\ell(g')$ is S_0 -minimal and, then, Lemma 2.12(iii) implies that $N_\ell(g')N_r(h)$ and $D_r(h)$ are right-coprime.

Now, let w_1, w_2 be words of $\mathcal{W}^+(S)$ representing the left denominator and numerator of g , let v_1, v_2 be words of $\mathcal{W}^+(S)$ representing the left denominator and numerator of g' , and let u_1, u_2 be words of $\mathcal{W}^+(S_0)$ representing the right numerator and denominator of h . Finally, let u represent f . First, by (the left counterpart of) Lemma 1.13, we have $w \overset{0,1,2}{\rightsquigarrow}_R w_1^{-1}w_2$. Next, $N_\ell(g) = fN_\ell(g')$ implies $w_1 \equiv_R^+ uv_1$, whence $w_1^{-1} \overset{1}{\rightsquigarrow}_R v_1^{-1}u^{-1}$ and $w \overset{0,1,2}{\rightsquigarrow}_R v_1^{-1}u^{-1}w_2$. Finally, we have $f^{-1}N_\ell(g) = N_\ell(g')N_r(h)D_r(h)^{-1}$, and we saw above that $N_\ell(g')N_r(h)$ and $D_r(h)$ are right-coprime, which implies that $N_\ell(g')N_r(h)$ is the right-numerator of $f^{-1}N_\ell(g)$ and $D_r(h)$ is its right-denominator. By definition, $u^{-1}w_2$ represents $f^{-1}N_\ell(g)$, whereas v_2u_1 represents $N_\ell(g')N_r(h)$ and u_2 represents $D_r(h)$. Lemma 1.13(ii) then implies $u^{-1}w_2 \overset{0,1,2}{\rightsquigarrow}_R v_2u_1u_2^{-1}$, whence $w \overset{0,1,2}{\rightsquigarrow}_R v_1^{-1}v_2u_1u_2^{-1}$, that is, $w \overset{0,1,2}{\rightsquigarrow}_R vu$ if we put $v = v_1^{-1}v_2$ and $u = u_1u_2^{-1}$. Now, by construction, v represents g' , an element of $\mathbf{T}(S_0)$, whereas u is a word of $\mathcal{W}(S_0)$. So (2.3) is satisfied. \square




FIGURE 5. Proof of Proposition 2.7: the point is that $D_\ell(g')$ and $N_\ell(g')N_r(h)$ are left-coprime, and that $N_\ell(g')N_r(h)$ and $D_r(h)$ are right-coprime.

2.3. Preservation under union. We shall now prove that Property H^\sharp is preserved under suitable unions.

Proposition 2.14. *Assume that (S_1, R_1) and (S_2, R_2) are positive group presentations that are in amalgam position and satisfy Property H^\sharp . Then $(S_1 \cup S_2, R_1 \cup R_2)$ satisfies Property H^\sharp .*

As explained in Subsection 2.1, this result together with Proposition 2.7 implies that every Artin–Tits presentation of FC type satisfies Property H^\sharp , hence a fortiori Property H, thus completing the proof of Proposition 2.1.

We split the proof of Proposition 2.14 into several steps. Until the end of the section, we assume that (S_1, R_1) and (S_2, R_2) are fixed positive presentations that are in amalgam position and satisfy Property H^\sharp . For $e = 1, 2$, we denote by G_e the group $\langle S_e | R_e \rangle$, we write $G_{1,2}$ for $G_1 \cap G_2$, and G for the amalgamated product $G_1 *_{G_{1,2}} G_2$. Thus, if we put $S = S_1 \cup S_2$, $R = R_1 \cup R_2$, and $S_{1,2} = S_1 \cap S_2$, then, by hypothesis, we have $G = \langle S | R \rangle$ and $G_{1,2} = \langle\langle S_{1,2} \rangle\rangle$ (both in the sense of G_1 and of G_2). Next, we fix an S_e -sequence of transversals \mathbf{T}_e witnessing that (S_e, R_e) satisfies Property H^\sharp , and write $\mathcal{P}_e(S', S'', w)$ for $\mathcal{P}_{\mathbf{T}_e}(S', S'', w)$.

For g in $G_1 \triangle G_2$ (that is, in $G_1 \setminus G_2$ or in $G_2 \setminus G_1$), we denote by $o(g)$ the index e such that g lies in G_e . It follows from Lemma 2.2 that, for every element g of G , there exists a unique sequence (g_1, \dots, g_r, g_*) such that $g = g_1 \cdots g_r g_*$ holds, for every i the element g_i is a nontrivial element of $\mathbf{T}_1(S_{1,2}) \cup \mathbf{T}_2(S_{1,2})$ with $o(g_i) \neq o(g_{i+1})$, and g_* is an element of $G_{1,2}$.

Definition 2.15. In the above context, the sequence (g_1, \dots, g_r, g_*) is called the *amalgam decomposition* of g and denoted by $\text{AD}(g)$. The number r is called the *amalgam length* of g and denoted by $\ell(g)$. Finally, the *tail* $\tau(g)$ of g is defined to be $g_r g_*$ for $r \geq 1$, and to be g_* (that is, g) for $r = 0$.

(Note that $\tau(g)$ is always $g_r g_*$ if $\text{AD}(g)$ is extended with $g_i = 1$ for $i \leq 0$.) We begin with a natural result saying that, if a word w represents an element g of G , then, by using special transformations, we can go from w to a concatenation of words that represent the successive entries of the amalgam decomposition of g . So, roughly speaking, the amalgam decomposition can be computed using special transformations.

Lemma 2.16. *Assume that w is a word of $\mathcal{W}(S)$ and (g_1, \dots, g_r, g_*) is the amalgam decomposition of \bar{w} . Then there exist words w_1, \dots, w_r, w_* such that $w \overset{0,1,2}{\rightsquigarrow}_R w_1 \cdots w_r w_*$ holds, w_i lies in $\mathcal{W}(S_{o(g_i)})$ and represents g_i for $i = 1, \dots, r$, and w_* lies in $\mathcal{W}(S_{1,2})$ and represents g_* .*

Proof. As S is the union of S_1 and S_2 , every word of $\mathcal{W}(S)$ can be expressed as a concatenation of words in $\mathcal{W}(S_1)$ and in $\mathcal{W}(S_2)$, say $w_1 \cdots w_p$ with $w_i \in \mathcal{W}(S_{e_i})$ and $e_i \neq e_{i+1}$. For w in $\mathcal{W}(S)$, we denote by $a(w)$ the minimal value of p for which such a decomposition of w exists. Then we prove the result using induction on $a(w)$.

Assume first $a(w) = 1$. Then w belongs to $\mathcal{W}(S_e)$ for $e = 1$ or 2 . Two subcases are possible. Assume first $\bar{w} \in G_{1,2}$. Then we have $r = 0$, with $\text{AD}(\bar{w}) = (g)$. By Lemma 2.6(ii), there exists w_* in $\mathcal{W}(S_{1,2})$ satisfying $w \overset{0,1,2}{\rightsquigarrow}_{R_1} w_*$. A fortiori, we have $w \overset{0,1,2}{\rightsquigarrow}_R w_*$, whence $\overline{w_*} = \bar{w}$, and $\overline{w_*} = g = g_*$, as expected. Assume now $\bar{w} \notin G_{1,2}$. Then, we have $r = 1$. By $\mathcal{P}_e(S_e, S_{1,2}, w)$, there exists w_1 in $\mathcal{W}(S_e)$ and w_* in $\mathcal{W}(S_{1,2})$ satisfying $w \overset{0,1,2}{\rightsquigarrow}_{R_e} w_1 w_*$ and $\overline{w_1} \in \mathbf{T}_e(S_{1,2})$. A fortiori, we have $w \overset{0,1,2}{\rightsquigarrow}_R w_1 w_*$, whence $\overline{w_1 w_*} = \bar{w}$. As, by construction, $\overline{w_*}$ belongs to $G_{1,2}$, the uniqueness of the amalgam decomposition implies $\overline{w_1} = g_1$ and $\overline{w_*} = g_*$, as expected again.

Assume now $a(w) \geq 2$. Write $w = vu$ with $a(v) = a(w) - 1$ and $a(u) = 1$. Then u belongs to $\mathcal{W}(S_e)$ for $e = 1$ or 2 . Let (f_1, \dots, f_q, f_*) be the amalgam decomposition of \bar{v} . By induction hypothesis, there exist words v_1, \dots, v_q, v_* such that $v \overset{0,1,2}{\rightsquigarrow}_R v_1 \cdots v_q v_*$ holds, v_i lies in $\mathcal{W}(S_{o(f_i)})$ and represents f_i for $i = 1, \dots, q$, and v_* lies in $\mathcal{W}(S_{1,2})$ and represents f_* . We consider the various ways $\text{AD}(\bar{w})$ can be obtained from $\text{AD}(\bar{v})$ and from \bar{u} , hereafter denoted by h , an element of G_e .

We consider $f_* h$, which, by hypothesis, lies in G_e , and its word representative $v_* u$, which lies in $\mathcal{W}(S_e)$ and satisfies $a(v_* u) = 1$. Assume first $h \in G_{1,2}$. In this case, $f_* h$ lies in $G_{1,2}$ as well, and we have $r = q$ with $\text{AD}(\bar{w}) = (f_1, \dots, f_q, f_* h)$. By induction hypothesis (or by Lemma 2.6(ii) applied in $\langle S_e | R_e \rangle$), there exist a word u_* in $\mathcal{W}(S_{1,2})$ such that $v_* u \overset{0,1,2}{\rightsquigarrow}_R u_*$ holds and, therefore, u_* represents $f_* u$. Then we have $w \overset{0,1,2}{\rightsquigarrow}_R v_1 \cdots v_q u_*$, and the words v_1, \dots, v_q, u_* have the expected properties.

Assume now $h \in G_e \setminus G_{1,2}$ with $q = 0$ or $o(f_q) \neq e$. In this case, $f_* h$ lies in $G_e \setminus G_{1,2}$, and we have $r = q + 1$ with $\text{AD}(\bar{w}) = (f_1, \dots, f_q, h_1, h_*)$ where (h_1, h_*) is the amalgam decomposition of $f_* h$. Now, as above, we have $a(v_* u) = 1$, so, by induction hypothesis, there exist words u_1, u_* such that $v_* u \overset{0,1,2}{\rightsquigarrow}_{R_e} u_1 u_*$ holds, u_1 lies in $\mathcal{W}(S_e)$ and represents h_1 , and u_* lies in $\mathcal{W}(S_{1,2})$ and represents h_* . Then we have $w \overset{0,1,2}{\rightsquigarrow}_R v_1 \cdots v_q v_* u \overset{0,1,2}{\rightsquigarrow}_R v_1 \cdots v_q u_1 u_*$, and the words $v_1, \dots, v_q, u_1, u_*$ have the expected properties.

Assume now $h \in G_e \setminus G_{1,2}$ with $q \geq 1$ and $o(f_q) = e$. Then we consider $f_q f_* h$, which, by hypothesis, lies in G_e , and its word representative $v_q v_* u$: by hypothesis, v_q lies in $\mathcal{W}(S_{o(f_q)})$, which is $\mathcal{W}(S_e)$, so $v_q v_* u$ lies in $\mathcal{W}(S_e)$ and we have $a(v_q v_* u) = 1$. As above in the case of $v_* u$, two cases arise. Assume first $f_q f_* h \in G_{1,2}$. In this case, we have $r = q - 1$ and $\text{AD}(g) = (f_1, \dots, f_{q-1}, f_q f_* h)$. By induction hypothesis (or by Lemma 2.6), there exists u_* of $\mathcal{W}(S_{1,2})$ satisfying $v_q v_* u \overset{0,1,2}{\rightsquigarrow}_R u_*$ and, therefore, u_* represents $f_q f_* u$. Then we have $w \overset{0,1,2}{\rightsquigarrow}_R v_1 \cdots v_q v_* u \overset{0,1,2}{\rightsquigarrow}_R v_1 \cdots v_{q-1} u_*$, and v_1, \dots, v_{q-1}, u_* have the expected properties.

Assume finally $f_q f_* h \in G_e \setminus G_{1,2}$. Then we have $r = q$ with $\text{AD}(g) = (f_1, \dots, f_{q-1}, h_1, h_*)$ where (h_1, h_*) is the amalgam decomposition of $f_q f_* h$. As above, we have $a(v_q v_* u) = 1$ so, by induction hypothesis, there exist u_1 and u_* such that $v_* u \overset{0,1,2}{\rightsquigarrow}_R u_1 u_*$ holds, u_1 lies in $\mathcal{W}(S_e)$ and represents h_1 , and u_* lies in $\mathcal{W}(S_{1,2})$ and represents h_* . Then we have $w \overset{0,1,2}{\rightsquigarrow}_R v_1 \cdots v_q v_* u \overset{0,1,2}{\rightsquigarrow}_R v_1 \cdots v_{q-1} u_1 u_*$, and $v_1, \dots, v_{q-1}, u_1, u_*$ have the expected properties. \square

Lemma 2.17. *Assume that S' is included in S and g lies in $\langle\langle S' \rangle\rangle$. Then all entries of the amalgam decomposition of g lie in $\langle\langle S' \rangle\rangle$. More precisely, if $\text{AD}(g)$ is (g_1, \dots, g_r, g_*) , then g_i lies in $\langle\langle S' \cap S_{o(g_i)} \rangle\rangle$ for $i = 1, \dots, r$, and g_* lies in $\langle\langle S' \cap S_{1,2} \rangle\rangle$.*

Proof. As S' is the union of $S' \cap S_1$ and $S' \cap S_2$, every element of $\langle\langle G' \rangle\rangle$ can be expressed as a product of elements of $\langle\langle S' \cap S_1 \rangle\rangle$ and $\langle\langle S' \cap S_2 \rangle\rangle$, say $g_1 \cdots g_p$ with $g_i \in \langle\langle S' \cap S_{e_i} \rangle\rangle$ and $e_i \neq e_{i+1}$. For g in $\langle\langle S' \rangle\rangle$, we define $\ell_{S'}(g)$ to be the minimal value of p for which such a decomposition of g exists. Then we prove the result using induction on $\ell_{S'}(g)$.

Assume first $\ell_{S'}(g) = 1$. Then g belongs to G_e for $e = 1$ or 2 , and $\ell(g) \leq 1$ holds. If g lies in $G_{1,2}$, the amalgam decomposition of g is (g) , and g lies in $\langle\langle S' \rangle\rangle$. Assume now $g \notin G_{1,2}$. Let w be a word of $\mathcal{W}(S)$ that represents g . By $\mathcal{P}_e(S' \cap S_e, S_{1,2}, w)$, we find w_1 in $\mathcal{W}(S' \cap S_e)$ and w_* in $\mathcal{W}(S' \cap S_{1,2})$ satisfying $w \overset{0,1,2}{\rightsquigarrow}_{R_e} w_1 w_*$ and $\overline{w_1} \in \mathbf{T}_e(S_{1,2})$. By uniqueness, the amalgam decomposition of g must be $(\overline{w_1}, \overline{w_*})$ and, by construction, $\overline{w_1}$ lies in $\langle\langle S' \cap S_e \rangle\rangle$ and $\overline{w_*}$ lies in $\langle\langle S' \cap S_{1,2} \rangle\rangle$.

Assume now $\ell_{S'}(g) \geq 2$. Write $g = fh$ with f, h in $\langle\langle S' \rangle\rangle$ satisfying $\ell_{S'}(f) < \ell_{S'}(g)$ and $\ell_{S'}(h) = 1$. Let (f_1, \dots, f_q, f_*) be the amalgam decomposition of f . By induction hypothesis, f_1, \dots, f_q and f_* lie in $\langle\langle S' \rangle\rangle$. Then, according to the scheme used in the proof of Lemma 2.16, the amalgam decomposition of g is of one of the types

- (i) $(f_1, \dots, f_q, f_* h)$,
- (ii) $(f_1, \dots, f_q, h_1, h_*)$ with $\ell_{S'}(f_* h) = 1$ and $(h_1, h_*) = \text{AD}(f_* h)$,
- (iii) $(f_1, \dots, f_{q-1}, f_q f_* h)$,
- (iv) $(f_1, \dots, f_{q-1}, h_1, h_*)$ with $\ell_{S'}(f_q f_* h) = 1$ and $(h_1, h_*) = \text{AD}(f_q f_* h)$.

By hypothesis, all involved elements, in particular h_1 and h_* in cases (ii) and (iv) owing to the induction hypothesis, lie in $\langle\langle S' \cap S_1 \rangle\rangle$, $\langle\langle S' \cap S_2 \rangle\rangle$, or $\langle\langle S' \cap S_{1,2} \rangle\rangle$ as expected. \square

Using Lemmas 2.17 and 2.6(ii), we can obtain a more precise version of Lemma 2.16.

Lemma 2.18. *Assume that S' is included in S and w is a word of $\mathcal{W}(S)$ that represents an element of $\langle\langle S' \rangle\rangle$. Let (g_1, \dots, g_r, g_*) be the amalgam decomposition of \overline{w} . Then there exist words w'_1, \dots, w'_r, w'_* such that $w \overset{0,1,2}{\rightsquigarrow}_R w'_1 \cdots w'_r w'_*$ holds, w'_i lies in $\mathcal{W}(S' \cap S_{o(g_i)})$ and represents g_i for $i = 1, \dots, r$, and w'_* lies in $\mathcal{W}(S' \cap S_{1,2})$ and represents g_* .*

The difference with Lemma 2.16 is that, here, we assume that the words w'_i and w'_* lie in $\mathcal{W}(S')$, that is, involve letters from S' only.

Proof. By Lemma 2.16, we first find w_1, \dots, w_r, w_* that represent g_1, \dots, g_r, g_* . Now, by Lemma 2.17, the elements g_1, \dots, g_r, g_* lie in $\langle\langle S' \rangle\rangle$. Then Lemma 2.6(ii) implies the existence of w'_1, \dots, w'_r, w'_* that satisfy $w_i \overset{0,1,2}{\rightsquigarrow}_R w'_i$ for $i = 1, \dots, r$ and $w_* \overset{0,1,2}{\rightsquigarrow}_R w'_*$, and all lie in $\mathcal{W}(S')$ in addition to the other properties. Then we have $w \overset{0,1,2}{\rightsquigarrow}_R w_1 \cdots w_r w_* \overset{0,1,2}{\rightsquigarrow}_R w'_1 \cdots w'_r w'_*$. \square

Our aim is to prove that (S, R) satisfies Property H^\sharp and, therefore, we must construct, for every subset S_0 of S , a distinguished $\langle\langle S_0 \rangle\rangle$ -transversal $\mathbf{T}(S_0)$ in the group G . To this end, we use the transversals $\mathbf{T}_1(S_0 \cap S_1)$ and $\mathbf{T}_2(S_0 \cap S_2)$. The natural idea is that $\mathbf{T}(S_0)$ consists of those elements of G whose tail belongs to $\mathbf{T}_e(S_0 \cap S_e)$ for $e = 1$ or 2 —we recall from Definition 2.15 that the tail $\tau(g)$ of g is the maximal final chunk of g that lies in G_1 or G_2 . To avoid the ambiguities resulting from the fact that, on $G_1 \cap G_2$, the transversals $\mathbf{T}_1(S_0 \cap S_1)$ and $\mathbf{T}_2(S_0 \cap S_2)$ need not coincide, we systematically choose $\mathbf{T}_1(S_0 \cap S_1)$ whenever possible.

Definition 2.19. For S_0 included in S , we put

$$(2.5) \quad \mathbf{T}(S_0) = \{g \in G \mid \tau(g) \in \mathbf{T}_1(S_0 \cap S_1) \cup (\mathbf{T}_2(S_0 \cap S_2) \setminus G_1)\}.$$

We shall show that \mathbf{T} is an S -sequence of transversals witnessing for Property H^\sharp . Note that, the hypothesis that $\mathbf{T}_1(S_0 \cap S_1)$ contains 1 implies that $\mathbf{T}(S_0)$ contains 1 as well. We begin with two preparatory results.

Lemma 2.20. *Assume that S_0 is included in S , that g lies in $\mathbf{T}(S_0)$ and that g' lies in the left- $\langle\langle S_0 \rangle\rangle$ -coset of g . Let (g_1, \dots, g_r, g_*) and $(g'_1, \dots, g'_{r'}, g'_*)$ be the amalgam decompositions of g and g' . Then we have (i) $r' \geq r$, (ii) $o(g'_r) = o(g_r)$ if $r \geq 1$ holds, and (iii) $g'_i = g_i$ for $1 \leq i < r$.*

Proof. By hypothesis, there exists h in $\langle\langle S_0 \rangle\rangle$ such that $g' = gh$ holds. Let (h_1, \dots, h_q, h_*) be the amalgam decomposition of h . By Lemma 2.17 applied to $\langle\langle S_0 \rangle\rangle$, the elements h_1, \dots, h_q, h_* all lie in $\langle\langle S_0 \rangle\rangle$. The results are obvious or vacuously true for $r = 0$, and we assume $r \geq 1$, with $o(g_r) = e$. We determine $\text{AD}(g')$ starting from $\text{AD}(g)$ and $\text{AD}(h)$ using arguments similar to those of Lemma 2.16. The point is that the case when the length of the amalgam decomposition decreases turns out to be impossible. Write \bar{e} for $3 - e$.

For $q = 0$, we find $\text{AD}(g') = (g_1, \dots, g_r, hh_*)$, and $\ell(g') = r$, so everything is clear.

Next assume $q \geq 1$ with $h_1 \in \mathbf{T}_{\bar{e}}(S_{1,2})$. Assume $\text{AD}(g_*h) = (f_1, \dots, f_p, f_*)$. By hypothesis, g_*h_1 lies in $G_{\bar{e}} \setminus G_{1,2}$, which, by uniqueness of the amalgam decomposition, implies $f_1 \in \mathbf{T}_{\bar{e}}(S_{1,2})$ (and $p = q$, but this is not needed here). In this case, we have $\text{AD}(gg_0) = (g_1, \dots, g_r, f_1, \dots, f_p, f_*)$. We deduce $\ell(g') = r + p \geq r$ and the other expected results.

Assume now $q \geq 1$ with $h_1 \in \mathbf{T}_e(S_{1,2})$ and $g_r g_* h_1 \notin G_{1,2}$. Let $\text{AD}(g_r g_* h) = (f_1, \dots, f_p, f_*)$. By hypothesis, $g_r g_* h_1$ lies in $G_e \setminus G_{1,2}$, which implies $p \geq 1$ and $f_1 \in \mathbf{T}_e(S_{1,2})$. In this case, we find $\text{AD}(g') = (g_1, \dots, g_{r-1}, f_1, \dots, f_p, f_*)$. We deduce $\ell(g') = r - 1 + p \geq r$ and the other expected results.

Assume finally $q \geq 1$ with $h_1 \in \mathbf{T}_e(S_{1,2})$ and $g_r g_* h_1 \in G_{1,2}$. By definition, the hypothesis that g belongs to $\mathbf{T}(S_0)$ implies that $g_r g_*$ lies in $\mathbf{T}_e(S_0 \cap S_e)$. On the other hand, by Lemma 2.17 applied with $S' = S_0$, the hypothesis that h lies in $\langle\langle S_0 \rangle\rangle$ implies that h_1 lies in $\langle\langle S_0 \cap S_e \rangle\rangle$. Here comes the point. Let w be any word of $\mathcal{W}(S_e)$ that represents $g_r g_* h_1$. Then $\mathcal{P}_e(S_{1,2}, S_0 \cap S_e, w)$ implies the existence of v in $\mathcal{W}(S_{1,2})$ and u in $\mathcal{W}(S_{1,2} \cap S_0)$ satisfying $w \overset{0,1,2}{\rightsquigarrow}_{R_e} vu$ and $\bar{v} \in \mathbf{T}_e(S_0 \cap S_e)$. Now, as h_1 lies in $\langle\langle S_0 \cap S_e \rangle\rangle$ and $\mathbf{T}(S_0 \cap S_e)$ is a $\langle\langle S_0 \cap S_e \rangle\rangle$ -transversal, we must have $\bar{v} = g_r g_*$ and $\bar{u} = h_1$. But this implies $g_r g_* \in G_{1,2}$, whence $g_r \in G_{1,2}$, which contradicts the definition of $\text{AD}(g)$. This case, which is the one that could possibly lead to $\ell(g') < \ell(g)$ is therefore impossible. So the proof is complete. \square

Lemma 2.21. *For every S_0 included in S , the set $\mathbf{T}(S_0)$ does not contain two distinct elements lying in the same left- $\langle\langle S_0 \rangle\rangle$ -coset.*

Proof. Assume that g, g' belong to $\mathbf{T}(S_0)$ and lie in the same left- $\langle\langle S_0 \rangle\rangle$ -coset of G . By Lemma 2.20, we have $\ell(g') \geq \ell(g)$, and $\ell(g) \geq \ell(g')$, whence $\ell(g) = \ell(g')$. Put $h = g^{-1}g'$, an element of $\langle\langle S_0 \rangle\rangle$. We now prove $g = g'$ by considering the various possible values of $\ell(g)$.

Assume first $\ell(g) = 0$. Then, by definition, g and g' belong to $\mathbf{T}_1(S_0 \cap S_1)$ (and to $G_{1,2}$), hence h belongs to $\langle\langle S_0 \rangle\rangle \cap G_1$, which, by Lemma 2.6, is $\langle\langle S_0 \cap S_1 \rangle\rangle$. So g and g' belong to the same left- $\langle\langle S_0 \cap S_1 \rangle\rangle$ -coset in G_1 . Then, the hypothesis that $\mathbf{T}_1(S_0 \cap S_1)$ is a $\langle\langle S_0 \cap S_1 \rangle\rangle$ -transversal in G_1 implies $g = g'$.

Assume now $\ell(g) = 1$. By definition, there exist e and e' such that g belongs to $\mathbf{T}_e(S_0 \cap S_e)$ and g' belongs to $\mathbf{T}_{e'}(S_0 \cap S_{e'})$. By Lemma 2.20(ii), we must have $e = e'$. It follows that h belongs to $\langle\langle S_0 \rangle\rangle \cap G_e$, which is $\langle\langle S_0 \cap S_e \rangle\rangle$. So g and g' belong to the same left- $\langle\langle S_0 \cap S_e \rangle\rangle$ -coset in G_e and, as $\mathbf{T}_e(S_0 \cap S_e)$ is a $\langle\langle S_0 \cap S_e \rangle\rangle$ -transversal in G_e , we must have $g = g'$.

Assume finally $r \geq 2$. By Lemma 2.20(iii), the amalgam decompositions of g and g' have the form $\text{AD}(g) = (g_1, \dots, g_r, g_*)$ and $\text{AD}(g') = (g_1, \dots, g_{r-1}, g'_r, g'_*)$. By definition of \mathbf{T} , the elements $\tau(g)$ and $\tau(g')$ belong to $\mathbf{T}(S_0)$ since g and g' do, and $g' = gh$ implies

$$\tau(g') = g'_r g'_* = g_1^{-1} \cdots g_{r-1}^{-1} g' = g_1^{-1} \cdots g_{r-1}^{-1} gh = g_r g_* h = \tau(g)h,$$

so $\tau(g')$ and $\tau(g)$ lie in the same $\langle\langle S_0 \rangle\rangle$ -coset. As, by construction, we have $\ell(\tau(g)) = \ell(\tau(g')) = 1$, the induction hypothesis implies $\tau(g) = \tau(g')$, whence $g = g'$. \square

We are now ready to complete the argument.

Proof of Proposition 2.14. Let S', S_0 be included in S . Two points have to be established, namely that $\mathbf{T}(S_0)$ is a $\langle\langle S_0 \rangle\rangle$ -transversal in G and that $\mathcal{P}_{\mathbf{T}}(S', S_0, w)$ holds for every word w of $\mathcal{W}(S)$ that represents an element of $\langle\langle S' \rangle\rangle$. We begin with the second point, which we establish using induction on $\ell(\bar{w})$.

(i) Assume first $\ell(\bar{w}) \leq 1$ with $\bar{w} \in G_1$. Applying Lemma 2.18 to S' and w , we first find w' in $\mathcal{W}(S' \cap S_1)$ satisfying $w \stackrel{0,1,2}{\rightsquigarrow}_R w'$. Then, applying $\mathcal{P}_1(S' \cap S_1, S' \cap S_0 \cap S_1, w')$, we find v in $\mathcal{W}(S' \cap S_1)$ and u in $\mathcal{W}(S' \cap S_0 \cap S_1)$ satisfying $w' \stackrel{0,1,2}{\rightsquigarrow}_{R_1} vu$ and $\bar{v} \in \mathbf{T}_1(S_0 \cap S_1)$. By composing, we deduce $w \stackrel{0,1,2}{\rightsquigarrow}_R vu$ and, by definition, $\mathbf{T}_1(S_0 \cap S_1)$ is included in $\mathbf{T}(S_0)$, so v and u witness for $\mathcal{P}_{\mathbf{T}}(S', S_0, w)$.

(ii) Assume now $\ell(\bar{w}) \leq 1$ with $\bar{w} \in G_2 \setminus G_1$. Arguing as above, but now in G_2 , we find v in $\mathcal{W}(S' \cap S_2)$ and u in $\mathcal{W}(S' \cap S_0 \cap S_2)$ satisfying $w \stackrel{0,1,2}{\rightsquigarrow}_{R_2} vu$ and $\bar{v} \in \mathbf{T}_2(S_0 \cap S_2)$. By definition, $\mathbf{T}_2(S_0 \cap S_2) \setminus G_1$ is included in $\mathbf{T}(S_0)$, so, if \bar{v} does not lie in G_1 , the words v and u witness for $\mathcal{P}_{\mathbf{T}}(S', S_0, w)$. If \bar{v} lies in G_1 , hence in $G_{1,2}$, one more step is needed to go from $\mathbf{T}_2(S_0 \cap S_2)$ to $\mathbf{T}_1(S_0 \cap S_1)$. Now, we have $\ell(\bar{u}) = 0$ with $\bar{u} \in G_1$, so by (i) we can find v' in $\mathcal{W}(S' \cap S_1)$ and u' in $\mathcal{W}(S' \cap S_0 \cap S_1)$ satisfying $v \stackrel{0,1,2}{\rightsquigarrow}_{R_1} v'u'$ and $\bar{v}' \in \mathbf{T}_1(S_0 \cap S_1)$, hence $\bar{v}' \in \mathbf{T}(S_0)$. Then we deduce $w \stackrel{0,1,2}{\rightsquigarrow}_R v'(u')$, and the words v' and $u'u$ witness for $\mathcal{P}_{\mathbf{T}}(S', S_0, w)$.

(iii) Assume now $\ell(\bar{w}) \geq 2$. Let $\text{AD}(\bar{w}) = (g_1, \dots, g_r, g_*)$. Applying Lemma 2.18 to S' and w , we find words w_1, \dots, w_r, w_* such that $w \stackrel{0,1,2}{\rightsquigarrow}_R w_1 \cdots w_r w_*$ holds, w_i lies in $\mathcal{W}(S' \cap S_{o(g_i)})$ and represents g_i for $i = 1, \dots, r$, and w_* lies in $\mathcal{W}(S' \cap S_{1,2})$ and represents g_* . Let $e = o(g_r)$. By construction, we have $\ell(g_r g_*) = 1$ with $g_r g_* \in G_e$. By (i) or (ii) applied to $w_r w_*$, we find words v_r in $\mathcal{W}(S' \cap S_e)$ and u in $\mathcal{W}(S' \cap S_0 \cap S_e)$ satisfying $w_r w_* \stackrel{0,1,2}{\rightsquigarrow}_{R_e} v_r u$ and $\bar{v}_r \in \mathbf{T}_e(S_0 \cap S_e)$. So, if we put $v = w_1 \cdots w_{r-1} v_r$, then $w \stackrel{0,1,2}{\rightsquigarrow}_R vu$ is true.

Two cases are possible. If \bar{v}_r does not belong to $G_{1,2}$, then $(g_1, \dots, g_{r-1}, \bar{v}_r)$ is the amalgam decomposition of \bar{v} , and \bar{v} lies in $\mathbf{T}(S_0)$. Then the words v and u witness for $\mathcal{P}_{\mathbf{T}}(S', S_0, w)$. On the other hand, if \bar{v}_r belongs to $G_{1,2}$, we have $\ell(\bar{v}) < \ell(\bar{w})$. Then, by induction hypothesis, we can find v' in $\mathcal{W}(S')$ and u' in $\mathcal{W}(S' \cap S_0)$ satisfying $v \stackrel{0,1,2}{\rightsquigarrow}_R v'u'$ and $\bar{v}' \in \mathbf{T}(S_0)$. Then we deduce $w \stackrel{0,1,2}{\rightsquigarrow}_R v'(u')$, and the words v' and $u'u$ witness for $\mathcal{P}_{\mathbf{T}}(S', S_0, w)$.

So, the proof of $\mathcal{P}_{\mathbf{T}}(S', S_0, w)$ is complete, and the only thing still to be proven is that $\mathbf{T}(S_0)$ is an $\langle\langle S_0 \rangle\rangle$ -transversal. This amounts to proving two facts, namely that two distinct elements of $\mathbf{T}(S_0)$ never lie in the same $\langle\langle S_0 \rangle\rangle$ -coset, which has been done in Lemma 2.21,

and that, for every g in G , there exists an element of $\mathbf{T}(S_0)$ is the left- $\langle\langle S_0 \rangle\rangle$ -coset of g . Now the latter point follows from $\mathcal{P}_{\mathbf{T}}(S, S_0, w)$. Indeed, let w be any word representing g . By $\mathcal{P}_{\mathbf{T}}(S, S_0, w)$, we find v in $\mathcal{W}(S)$ and u in $\mathcal{W}(S_0)$ satisfying $w \overset{0,1,2}{\rightsquigarrow}_R vu$ and $\bar{v} \in \mathbf{T}(S_0)$. Then \bar{u} lies in $\langle\langle S_0 \rangle\rangle$, and \bar{v} is an element of $\mathbf{T}(S_0)$ lying in the left- $\langle\langle S_0 \rangle\rangle$ -coset of g . \square

Thus the proof that all Artin–Tits presentations of type FC satisfy Properties H[#] and H (Proposition 2.1) is complete.

3. A SYNTACTIC APPROACH

Our proof of Property H for type FC is essentially based on the existence of a distinguished normal form for the elements of the considered Artin–Tits group. Although the approach is simple, it seems unlikely that it could lead to a proof of Conjecture 1.6 as the construction of a similar normal form in the case of general Artin–Tits presentations seems out of reach. Also, the possible interest of Property H is to provide some control of a group even in the case when no solution to the word problem is known, and using a preexisting solution to that word problem to establish Property H is not natural. In this section, we develop another approach, which leads to Property H directly and does not appeal to any normal form or, more generally, to any solution of the word problem. The principle is to start from a derivation that involves transformations of type 0, 1, and ∞ and eliminate type ∞ transformations by introducing type 2 transformations instead.

We show in Subsection 3.1 how to realize this strategy in the case of right-angled Artin–Tits presentations. A few additional comments are included in Subsection 3.2.

3.1. Right-angled Artin–Tits presentations. An Artin–Tits presentation is said to be *right-angled* if it only involves commutation relations $st = ts$. Right-angled Artin–Tits presentations all are of FC type, so Proposition 2.5 implies

Proposition 3.1. *Every right-angled Artin–Tits presentation satisfies Property H.*

We shall now give a new proof of this result relying on a purely syntactic approach. The argument requires several auxiliary developments and will be completed at the end of the subsection. Until there, we assume that (S, R) is a fixed right-angled Artin–Tits presentation, and denote by G the group $\langle S | R \rangle$.

Assume that w is a word of $\mathcal{W}(S)$ that represents 1 in G . Then we have $w \overset{0,1,\infty}{\rightsquigarrow}_R \varepsilon$, so there exists a sequence of words w_0, \dots, w_m with $w_0 = w$, $w_m = \varepsilon$ and, for every k , the word w_k is obtained from w_{k-1} by applying a special transformation of type 0, 1, or ∞ . The idea will be to mark the additional letters created in type ∞ steps and to follow their subsequent destiny in the derivation. The point is that, when we erase these additional letters in a convenient way, the remaining words will make a derivation in terms of type 0, 1, and 2 transformations.

Implementing the above simple principle requires taking into account the step at which each additional letter is created. This leads us to introducing an augmented alphabet \underline{S} where all letters include a number that corresponds to their age, that is, the date of their creation and, accordingly, an augmented version \rightsquigarrow_R of the relation \rightsquigarrow_R for words of $\mathcal{W}(\underline{S})$.

Definition 3.2. We denote by \underline{S} the alphabet consisting of one letter $s_{[i]}$ for each letter s of S and each nonnegative integer i ; we identify s with $s_{[0]}$. We denote by ϕ the alphabetical homomorphism of $\mathcal{W}(\underline{S})$ to $\mathcal{W}(S)$ that maps every letter $s_{[i]}$ to s . For $h \geq 0$, we denote by π_h the alphabetical homomorphism of $\mathcal{W}(\underline{S})$ to itself that maps $s_{[i]}$ to itself for $i < h$ and to ε for $i \geq h$.

Words of $\mathcal{W}(\underline{S})$ will be called *augmented*. Thus, assuming that $\mathbf{a}, \mathbf{b}, \mathbf{c}$ are letters of S , a typical augmented word is $\mathbf{a}_1\mathbf{B}_0\mathbf{c}_2\mathbf{A}_1\mathbf{c}_0$. Its image under ϕ is the (standard) word \mathbf{aBcAc} (all indices are forgotten), whereas its image under π_2 is $\mathbf{a}_1\mathbf{B}_0\mathbf{A}_1\mathbf{c}_0$ (all letters with index 2 and larger are erased), and its image under π_1 is $\mathbf{B}_0\mathbf{c}_0$.

Definition 3.3. *Special transformations on augmented words are defined as follows:*

- type 0: Remove some pair $s_{[i]}^e s_{[j]}^{-e}$, $e = \pm 1$, and replace each remaining occurrence of $s_{[i]}^{\pm 1}$ and $s_{[j]}^{\pm 1}$ with $s_{[\min(i,j)]}^{\pm 1}$;
- type 1: Replace $s_{[i]}^e t_{[j]}^e$ with $t_{[j]}^e s_{[i]}^e$, where $st = ts$ is a relation of R and e is ± 1 ;
- type 2: Replace $s_{[i]}^e t_{[j]}^{-e}$ with $t_{[j]}^{-e} s_{[i]}^e$, where $st = ts$ is a relation of R and e is ± 1 ;
- type ∞ : Insert some pair $s_{[i]}^e s_{[i]}^{-e}$, $e = \pm 1$, where i is strictly larger than every index of letter occurring in the current word.

We write $w \xrightarrow{X}_R w'$ if w' can be obtained from w by a finite sequence of special transformations of type belonging to X .

Special transformations on augmented words are liftings of the corresponding special transformations on words: a straightforward inspection shows that, if an augmented word w' is the image of another augmented word w under a transformation of type i , then the (non-augmented) word $\phi(w')$ is the image of $\phi(w)$ under a transformation of type i .

The first observation is that every sequence of special transformations on standard words can be lifted into a sequence of special transformations on augmented words. Hereafter, we say that (w_0, w_1, \dots) is a *special sequence* of (possibly augmented) words if every word w_k is obtained from w_{k-1} using a special transformation.

Lemma 3.4. *For every special sequence (w_0, w_1, \dots) in $\mathcal{W}(S)$, there exists a special sequence of augmented words $(\widehat{w}_0, \widehat{w}_1, \dots)$ such that \widehat{w}_0 equals w_0 and, for every k , the word w_k is the image of \widehat{w}_k under ϕ and, moreover, the type of the transformation connecting \widehat{w}_{k-1} to \widehat{w}_k is the same as the type of the transformation connecting w_{k-1} to w_k .*

Proof. We start from $\widehat{w}_0 = w_0$ and use an induction. For steps of type 0, 1, or 2, we just mimick the transformation. For steps of type ∞ , that is, when w_k is obtained from w_{k-1} by inserting a pair $s^e s^{-e}$, we define \widehat{w}_k from \widehat{w}_{k-1} by inserting $s_{[i]}^e s_{[i]}^{-e}$, where i is the maximum of all indices of letters occurring in \widehat{w}_{k-1} augmented by 1. \square

Example 3.5. Assume $S = \{\mathbf{a}, \mathbf{b}, \mathbf{c}\}$ and let $R = \{\mathbf{ab} = \mathbf{ba}, \mathbf{bc} = \mathbf{cb}, \mathbf{ac} = \mathbf{ca}\}$. Then (S, R) is a right-angled Artin-Tits presentation. As in Section 1, consider $w = \mathbf{aBcAbC}$. Then, starting from the special sequence

$$\mathbf{aBcAbC} \xrightarrow{\infty} \mathbf{aBAcAbC} \xrightarrow{1}_R \mathbf{aABacAbC} \xrightarrow{0} \mathbf{BacAbC} \xrightarrow{1}_R \mathbf{BcaAbC} \xrightarrow{0} \mathbf{BcbC} \xrightarrow{1}_R \mathbf{Bbcc} \xrightarrow{0} \mathbf{cC} \xrightarrow{0} \varepsilon$$

and using the method of Lemma 3.4 gives

$$\begin{aligned} \mathbf{a}_0\mathbf{B}_0\mathbf{c}_0\mathbf{A}_0\mathbf{b}_0\mathbf{C}_0 &\xrightarrow{\infty} \mathbf{a}_0\mathbf{B}_0\mathbf{A}_1\mathbf{a}_1\mathbf{c}_0\mathbf{A}_0\mathbf{b}_0\mathbf{C}_0 \xrightarrow{1}_R \mathbf{a}_0\mathbf{A}_1\mathbf{B}_0\mathbf{a}_1\mathbf{c}_0\mathbf{A}_0\mathbf{b}_0\mathbf{C}_0 \\ &\xrightarrow{0} \mathbf{B}_0\mathbf{a}_0\mathbf{c}_0\mathbf{A}_0\mathbf{b}_0\mathbf{C}_0 \xrightarrow{1}_R \mathbf{B}_0\mathbf{c}_0\mathbf{a}_0\mathbf{A}_0\mathbf{b}_0\mathbf{C}_0 \xrightarrow{0} \mathbf{B}_0\mathbf{c}_0\mathbf{b}_0\mathbf{C}_0 \xrightarrow{1}_R \mathbf{B}_0\mathbf{b}_0\mathbf{c}_0\mathbf{C}_0 \xrightarrow{0} \mathbf{c}_0\mathbf{C}_0 \xrightarrow{0} \varepsilon, \end{aligned}$$

where we write \mathbf{a}_i for $\mathbf{a}_{[i]}$.

The main point now is to control the properties of the augmented words that appear when special sequences are lifted.

Definition 3.6. (i) An augmented word of the form $r_{[h]}^e w r_{[h]}^{-e}$ is called an *h-pair*; it is said to be *commutative* if $rs = sr$ is a relation of R for every letter s such that $s^{\pm 1}$ occurs in $\phi(\pi_h(w))$ (that is, such that some letter $s_{[i]}^{\pm 1}$ with $i \leq h$ occurs in w).

(ii) An augmented word w is said to be *regular* if, for every $h \geq 1$, either w contains no letter of index h , or it contains exactly two such letters forming a commutative h -pair.

The argument for Proposition 3.1 will be inductive. Here is the basic step for the induction.

Lemma 3.7. *Assume that w is a regular augmented word and w' is obtained from w using one special transformation. Then*

- (i) *The word w' is regular;*
- (ii) *If w or w' contains at least one letter of positive index and if h is the highest index of a letter occurring in w and w' , the relation $\pi_h(w) \overset{0,1,2,\infty}{\rightsquigarrow_R} \pi_h(w')$ holds;*
- (iii) *The relation $\pi_1(w) \overset{0,1,2}{\rightsquigarrow_R} \pi_1(w')$ holds.*

Proof. We first establish (i) and (ii) by considering the possible types of special transformations connecting w to w' . We begin with type 0, that is, we assume that w' is obtained from w by removing some factor $s_{[i]}^e s_{[j]}^{-e}$, $e = \pm 1$, and replacing the remaining occurrences of $s_{[i]}^{\pm 1}$ and $s_{[j]}^{\pm 1}$ with $s_{[\min(i,j)]}^{\pm 1}$.

(i) If h is neither i nor j , the possible letters of index h in w' are those of w , and the commutativity condition is unchanged (if the deleted factor is not nested in the possible h -pair), or weakened. In any case, the hypothesis that w satisfies the properties implies that w' does too. Assume now $h = i < j$. Then we must be in the case when w contains a commutative h -pair, say $s_{[h]}^e v s_{[h]}^{-e}$, and w' is obtained by removing one of the letters $s_{[h]}^{-e}$ together with some adjacent $s_{[j]}^{\pm 1}$, which occurs in some j -pair $s_{[j]}^d v' s_{[j]}^{-d}$. So, up to symmetries, we may assume that we are in the situation

$$\dots s_{[h]}^e v s_{[h]}^{-e} s_{[j]}^e v' s_{[j]}^{-e} \dots \overset{0}{\rightsquigarrow} \dots s_{[h]}^e v v' s_{[h]}^{-e} \dots,$$

and we see that the expected properties hold in w' : indeed, w' contains one h -pair, no j -pair, and the h -pair is commutative, because, by hypothesis, the letter s commutes with all letters of $\phi(\pi_h(v))$ and all letters of $\phi(\pi_j(v'))$, hence, a fortiori, with all letters of $\phi(\pi_h(v v'))$. The other cases are similar. Finally, assume $h = i > j$. Then the typical case is

$$\dots s_{[h]}^e v s_{[h]}^{-e} s_{[j]}^e v' s_{[j]}^{-e} \dots \overset{0}{\rightsquigarrow} \dots s_{[j]}^e v v' s_{[j]}^{-e} \dots,$$

and, again, the hypothesis that s commutes with all letters of $\phi(\pi_h(v))$ and all letters of $\phi(\pi_j(v'))$ implies that s commutes with all letters of $\phi(\pi_j(v v'))$. So w' is regular.

(ii) If i and j are strictly smaller than h , the letters of index i and j are invariant when π_h is applied, and the word $\pi_h(w')$ is obtained from $\pi_h(w)$ by a type 0 transformation, that is, $\pi_h(w) \overset{0}{\rightsquigarrow_R} \pi_h(w')$ holds. Assume now $i = h > j$. Up to symmetries, we are in the situation of

$$\dots s_{[h]}^e v s_{[h]}^{-e} s_{[j]}^e v' s_{[j]}^{-e} \dots \overset{0}{\rightsquigarrow} \dots s_{[j]}^e v v' s_{[j]}^{-e} \dots$$

When we apply π_h , the above words respectively become

$$\dots \pi_h(v) s_{[j]}^e \pi_h(v') s_{[j]}^{-e} \dots \quad \text{and} \quad \dots s_{[j]}^e \pi_h(v) \pi_h(v') s_{[j]}^{-e} \dots$$

The point is that, as, by hypothesis, the h -pair $s_{[h]}^e v s_{[h]}^{-e}$ is commutative, the letter s commutes with every letter occurring in $\pi_h(v)$, and, therefore, we have $\pi_h(v) s_{[j]}^e \overset{1,2}{\rightsquigarrow_R} s_{[j]}^e \pi_h(v)$, whence $\pi_h(w) \overset{1,2}{\rightsquigarrow_R} \pi_h(w')$.

We turn to type 1 and assume now that w' is obtained from w by replacing $s_{[i]}^e t_{[j]}^e$ with $t_{[j]}^e s_{[i]}^e$, where $st = ts$ is a relation of R and e is ± 1 .

(i) Here the same letters occur in w and w' , so the possible letters of index h occurring in w and w' are the same. In particular, there exists an h -pair in w if and only if there is one in w' . Moreover, the hypothesis that the possible h -pair is commutative in w implies that the corresponding h -pair in w' is commutative as well. Indeed, if h is neither i nor j ,

the letters nested in the h -pair are not changed; otherwise, up to symmetries, the typical case is

$$\dots s_{[i]}^e r_{[h]}^e v r_{[h]}^{-e} \dots \xrightarrow{\sim_R^1} \dots r_{[h]}^e s_{[i]}^e v r_{[h]}^{-e} \dots,$$

and the point is that the hypotheses that r commutes with $\phi(\pi_h(v))$ and, on the other hand, with s , implies that it commutes with $\phi(\pi_h(s_{[i]}^e v))$ in every case. So the possible h -pair of w' is commutative, and w' is regular

(ii) Again, if h is larger than i and j , removing the letters of index h changes nothing essential and we obtain $\pi_h(w) \xrightarrow{\sim_R^1} \pi_h(w')$. On the other hand, if $h = i > j$ holds, removing the letters of index h cancels the effect of the transformation, and we simply have $\pi_h(w) = \pi_h(w')$, hence (trivially) $\pi_h(w) \xrightarrow{\sim_R^{0,1,2,\infty}} \pi_h(w')$.

We now consider type 2 and assume that w' is obtained from w by replacing $s_{[i]}^e t_{[j]}^{-e}$ with $t_{[j]}^{-e} s_{[i]}^e$, where $st = ts$ is a relation of R and e is ± 1 . Both for (i) and (ii), the argument is similar to that of type 1, the only difference being that the commuted letters have now opposite signs (a specificity of the right-angled case).

Finally we consider type ∞ , that is, assume that w' is obtained from w by inserting some pair $s_{[i]}^e s_{[i]}^{-e}$, $e = \pm 1$, where i is strictly larger than every index of letter occurring in w .

(i) If h is not i , in which case $h < i$ holds by definition, the possible h -pair of w provides an h -pair in w' . We have no commutation hypothesis involving the letter s , but, even if the inserted pair is nested in the h -pair as in

$$\dots r_{[h]}^e v v' r_{[h]}^{-e} \dots \xrightarrow{\infty} \dots r_{[h]}^e v s_{[i]}^e s_{[i]}^{-e} v' r_{[h]}^{-e} \dots,$$

we are sure that r commutes with every letter of $\phi(\pi_h(v s_{[i]}^e s_{[i]}^{-e} v'))$ as the latter word coincides with $\phi(\pi_h(v v'))$. So w' is regular.

(ii) By definition, we have $\pi_h(w) = \pi_h(w')$, whence again $\pi_h(w) \xrightarrow{\sim_R^{0,1,2,\infty}} \pi_h(w')$.

So the proof of (i) and (ii) is complete, and it remains to deduce (iii). First, we claim that $\pi_h(w) \xrightarrow{\sim_R^{0,1,2,\infty}} \pi_h(w')$ holds for every $h \geq 1$. Indeed, the result is obvious if h is larger than every index of letter occurring in w and w' since, in this case, we have $\pi_h(w) = w$ and $\pi_h(w') = w'$. Then the result follows from (ii), using a decreasing induction on h since applying π_h to a word where the largest index is h yields a word where the largest index is at most $h - 1$. So we obtain $\pi_1(w) \xrightarrow{\sim_R^{0,1,2,\infty}} \pi_1(w')$. Now, the word $\pi_1(w')$ contains no letter of positive index. As no type ∞ transformation can lead to a word containing no letter with positive index, the only possibility is that $\pi_1(w) \xrightarrow{\sim_R^{0,1,2}} \pi_1(w')$ holds, as expected. \square

We can now combine the results to obtain the expected new proof of Proposition 3.1.

Proof of Proposition 3.1. Let w be a word of $\mathcal{W}(S)$ that represents 1 in $\langle S|R \rangle$. Then, by (1.1), we have $w \xrightarrow{\sim_R^{0,1,\infty}} \varepsilon$. Fix a special sequence w_0, w_1, \dots, w_m going from w to the empty word.

By Lemma 3.4, we can find a special sequence of augmented words $\widehat{w}_0, \widehat{w}_1, \dots$ such that \widehat{w}_0 coincides with w_0 and, for every k , the word w_k is the image of \widehat{w}_k under ϕ .

By definition, the (augmented) word \widehat{w}_0 contains no letter of positive index, hence it is regular. Hence, by Lemma 3.7(i), every augmented word \widehat{w}_k is regular. Then Lemma 3.7(iii) implies $\pi_1(\widehat{w}_{k-1}) \xrightarrow{\sim_R^{0,1,2}} \pi_1(\widehat{w}_k)$ for each k . We deduce $\pi_1(\widehat{w}_0) \xrightarrow{\sim_R^{0,1,2}} \pi_1(\widehat{w}_m)$, which is the expected relation $w \xrightarrow{\sim_R^{0,1,2}} \varepsilon$. \square

Example 3.8. Applying the above argument to the case of Example 3.5 amounts to collapsing all letters with positive index and forgetting indices. Starting from the initial sequence

that contains a type ∞ step (top), we lift it into an augmented sequence (middle, we drop the zero indices), and project it into a new sequence that contains no type ∞ steps (down):

$$\begin{array}{cccccccccccc}
\mathbf{aBcAbC} & \overset{\infty}{\rightsquigarrow} & \mathbf{aBAacAbC} & \overset{1}{\rightsquigarrow}_R & \mathbf{aABacAbC} & \overset{0}{\rightsquigarrow} & \mathbf{BacAbC} & \overset{1}{\rightsquigarrow}_R & \mathbf{BcaAbC} & \overset{0}{\rightsquigarrow} & \mathbf{BcbC} & \overset{1}{\rightsquigarrow}_R & \mathbf{BbccC} & \overset{0}{\rightsquigarrow} & \mathbf{cC} & \overset{0}{\rightsquigarrow} & \mathbf{\varepsilon} \\
\uparrow \phi & & \uparrow \phi \\
\mathbf{aBcAbC} & \overset{\infty}{\rightsquigarrow} & \mathbf{aBA_1a_1cAbC} & \overset{1}{\rightsquigarrow}_R & \mathbf{aA_1Ba_1cAbC} & \overset{0}{\rightsquigarrow} & \mathbf{BacAbC} & \overset{1}{\rightsquigarrow}_R & \mathbf{BcaAbC} & \overset{0}{\rightsquigarrow} & \mathbf{BcbC} & \overset{1}{\rightsquigarrow}_R & \mathbf{BbccC} & \overset{0}{\rightsquigarrow} & \mathbf{cC} & \overset{0}{\rightsquigarrow} & \mathbf{\varepsilon} \\
\downarrow \pi_1 & & \downarrow \pi_1 \\
\mathbf{aBcAbC} & = & \mathbf{aBcAbC} & = & \mathbf{aBcAbC} & \overset{2}{\rightsquigarrow}_R & \mathbf{BacAbC} & \overset{1}{\rightsquigarrow}_R & \mathbf{BcaAbC} & \overset{0}{\rightsquigarrow} & \mathbf{BcbC} & \overset{1}{\rightsquigarrow}_R & \mathbf{BbccC} & \overset{0}{\rightsquigarrow} & \mathbf{cC} & \overset{0}{\rightsquigarrow} & \mathbf{\varepsilon}.
\end{array}$$

(Of course, this example is nearly trivial as one type ∞ transformation only occurs in the initial sequence.)

3.2. Further remarks. Extending the method of Subsection 3.1 to more general Artin–Tits presentations is a challenge we leave open. The technical problem is to find augmented versions of type 2 transformations: in the right-angled case, the latter just are commutations, and the definition is clear; in more complicated cases, things are not so clear. Let us just mention that, in the case of a length 3 relation $sts = tst$, geometric interpretations suggest that the proper way of defining an augmented version of the type 2 transformation $s^{-1}t \overset{2}{\rightsquigarrow}_R tst^{-1}s^{-1}$ could be to put $s_{[i]}^{-1}t_{[j]} \overset{2}{\rightsquigarrow}_R t_{[k]}s_{[j]}t_{[i]}^{-1}s_{[k]}^{-1}$, where k is larger than every index of letter occurring in the current word.

Let us still mention two more approaches. First, a natural way for proving Conjecture 1.6 could be to establish the confluence of the relation $\overset{0,1,2}{\rightsquigarrow}_R$, namely to prove that, if we have both $w \overset{0,1,2}{\rightsquigarrow}_R w'$ and $w \overset{0,1,2}{\rightsquigarrow}_R w''$, then there exists a word w''' satisfying $w' \overset{0,1,2}{\rightsquigarrow}_R w'''$ and $w'' \overset{0,1,2}{\rightsquigarrow}_R w'''$. However, as the relation $\overset{0,1,2}{\rightsquigarrow}_R$ need not be Noetherian in general (that is, there may exist infinite special sequences), establishing confluence is not easy.

On the other hand, working with $\overset{0,1,2}{\rightsquigarrow}_R$ is in general difficult because this relation is highly non-deterministic: for every word w , there exist in general many words w' such that $w \overset{0,1,2}{\rightsquigarrow}_R w'$ is satisfied. In [10], an algorithm called handle reduction is constructed in the case of classical braids, that is, of Artin–Tits groups of type A. This algorithm, called *handle reduction*, extends free group reduction, and it returns for every word w an equivalent word $\text{HR}(w)$ with the property that, if w represents 1, then $\text{HR}(w)$ is the empty word. However, this algorithm is not a normal form, as $w \equiv_R w'$ need not imply $\text{HR}(w) = \text{HR}(w')$. A precise analysis of the algorithm shows that $w \overset{0,1,2}{\rightsquigarrow}_R \text{HR}(w)$ always holds and, therefore, we obtain another proof of Property H in this case. In this approach, handle reduction is used as a strategy for choosing, for every word w , a distinguished word among the many words w' that satisfy $w \overset{0,1,2}{\rightsquigarrow}_R w'$. It is natural to wonder whether such an approach, which a priori is less ambitious than constructing a normal form, might apply to more general situations.

REFERENCES

- [1] J.M. Alonso and al., Group theory from a geometrical viewpoint, E. Ghys, A. Haefliger, A. Verjovsky (eds); World Sci. Publ., River Edge, NJ (1991).
- [2] J.A. Altobelli, *The word problem for Artin groups of FC type*, J. Pure Appl. Algebra **129** (1998) 1–22.
- [3] K.I. Appel and P.E. Schupp, *Artin groups and infinite Coxeter groups*, Invent. Math. **72** (1983) 201–220.
- [4] E. Brieskorn, *Sur les groupe de tresses*, Séminaire Bourbaki 1971/72, Lectures Notes in Math. Springer-Verlag **317** (1973) 21–44.
- [5] E. Brieskorn and K. Saito, *Artin-Gruppen und Coxeter-Gruppen*, Invent. Math. **17** (1972) 245–271.
- [6] J. Cannon, *The theory of negative curved spaces and groups in Ergodic Theory, Symbolic Dynamics and Hyperbolic spaces*, T. Bedford, M. Keane, C. Series (eds); Oxford University Press (1991) 315–369.
- [7] R. Charney, *Geodesic automation and growth functions for Artin groups of finite type*, Math. Ann. **301** (1995) 307–324.

- [8] Charney, R., Problems related to Artin Groups, American Institute of Mathematics, Problems in Geometric Group Theory wiki, http://people.brandeis.edu/~charney/papers/Artin_probs3.pdf, <http://aimath.org/pggt>
- [9] P. Dehornoy, *Groups with a complemented presentation*, J. Pure Appl. Algebra **116** (1997) 115–137.
- [10] P. Dehornoy, *A fast method for comparing braids*, Advances in Math. **125** (1997) 200–235.
- [11] P. Dehornoy, *Complete positive group presentations*, J. Algebra **268** (2003) 156–197.
- [12] P. Dehornoy, *The subword reversing method*, Intern. J. Alg. and Comput. **21** (2011) 71–118.
- [13] P. Dehornoy, with I. Dynnikov, D. Rolfsen, B. Wiest, *Ordering Braids*, Mathematical Surveys and Monographs vol. 148, Amer. Math. Soc. (2008).
- [14] P. Dehornoy and B. Wiest, *On word reversing in braid groups*, Int. J. Algebra Comput. **16** (2006) 931–947.
- [15] P. Deligne, *Les immeubles des groupes de tresses généralisés*, Invent. Math. **17** (1972) 273–302.
- [16] D. Epstein, with J. Cannon, D. Holt, S. Levy, M. Paterson, and W. Thurston, *Word Processing in Groups*, Jones & Bartlett Publ. (1992).
- [17] F.A. Garside, *The braid group and other groups*, Quart. J. Math. Oxford **20-78** (1969) 235–254.
- [18] J.E. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge Univ. Texts (1989).
- [19] I. Kapovich and P. Schupp, *Relative hyperbolicity and Artin Groups*, Geom. Dedic. **107** (2004) 153–167.
- [20] D. Krammer, *Braid groups are linear*, Ann. Math. **151-1** (2002) 131–156.
- [21] I.G. Lysionok, *On some algorithmic properties of hyperbolic groups*, math. USSR Izv **35** (1990) 145–163.
- [22] L. Paris, *Artin monoids inject in their groups*, Comment. Math. Helv. **77** (2002) 609–637.
- [23] K. Tatsuoka, *An isoperimetric inequality for Artin groups of finite type*, Trans. Amer. Math. Soc. **339-2** (1993) 537–551.
- [24] H. van der Lek, *The homotopy type of complex hyperplane complements*, PhD thesis Nijmegen (1983).

LABORATOIRE DE MATHÉMATIQUES NICOLAS ORESME, UMR 6139, UCBN ET CNRS, 14032 CAEN, FRANCE

E-mail address: `patrick.dehornoy@unicaen.fr`

URL: `//www.math.unicaen.fr/~dehornoy`

LABORATOIRE DE MATHÉMATIQUES NICOLAS ORESME, UMR 6139, UCBN ET CNRS, 14032 CAEN, FRANCE

E-mail address: `eddy.godelle@unicaen.fr`

URL: `//www.math.unicaen.fr/~godelle`